arXiv:math/0702436v5 [math.NT] 4 Jun 2010

Calculation of ℓ-adic Local Fourier Transformations *

Lei Fu
Chern Institute of Mathematics and LPMC, Nankai University, Tianjin 300071, P. R. China
leifu@nankai.edu.cn

Abstract

We calculate the local Fourier transformations for a class of $\overline{\mathbb{Q}}_\ell$-sheaves. In particular, we verify a conjecture of Laumon and Malgrange ([18] 2.6.3). As an application, we calculate the local monodromy of ℓ-adic hypergeometric sheaves introduced by Katz ([15]). We also discuss the characteristic p analogue of the Turrittin-Levelt Theorem for D-modules.

Key words: Fourier transformation, local Fourier transformation, hypergeometric sheaf.

Mathematics Subject Classification: 14F20, 14G15, 11L05.

Introduction

The global ℓ-adic Fourier transformation was first introduced by Deligne. To study the local behavior of the global Fourier transformation, Laumon [18] discovered the stationary phase principle and introduced local Fourier transformations. All these transformations are defined by cohomological functors and are rarely computable. However, in [18] 2.6.3, Laumon and Malgrange give conjectural formulas of local Fourier transformations for a class of $\overline{\mathbb{Q}}_\ell$-sheaves. In this paper we calculate local Fourier transformations for a more general class of $\overline{\mathbb{Q}}_\ell$-sheaves. In particular, we prove the conjecture of Laumon and Malgrange. It turns out that to get the correct result, the conjectural formulas of Laumon and Malgrange have to be slightly modified. As an application of our result, we calculate the local monodromy of ℓ-adic hypergeometric sheaves introduced by Katz ([15]). The method used in this paper can also be used to show that the Fourier transformation of an analytic sheaf with meromorphic ramification defined in [19] still has meromorphic ramification, thus answering positively a question proposed by Ramero.

Throughout this paper, for any ring A, we use the notations

$$\mathbb{A}^1_A = \operatorname{Spec} A[t],\ \mathbb{G}_{m,A} = \operatorname{Spec} A[t, 1/t],\ \mathbb{P}^1_A = \operatorname{Proj} A[t_0, t_1].$$

*I would like to thank anonymous referees of various versions of the paper for communicating to me their deep insight on the stationary phase principle and the Legendre transformation, and for their many suggestions on improving the exposition of the paper. I also thank P. Deligne, L. Illusie and N. Katz for their help while preparing this paper. The research is supported by the NSFC (10525107).

Let k be a perfect field of characteristic p, $\bar{k}$ an algebraic closure of k, q any power of p, $\mathbb{F}_q$ the finite field with q elements contained in $\bar{k}$, and ℓ a prime number distinct from p. Fix a nontrivial additive character $\psi : \mathbb{F}_p \to \overline{\mathbb{Q}}_\ell^*$. The k-morphism

$$\wp : \mathbb{A}^1_k \to \mathbb{A}^1_k, \quad t \mapsto t^p - t$$

is a finite Galois étale covering space, and it defines an $\mathbb{F}_p$-torsor

$$0 \to \mathbb{F}_p \to \mathbb{A}^1_k \xrightarrow{\wp} \mathbb{A}^1_k \to 0.$$

Pushing-forward this torsor by ψ^{-1}, we get a lisse $\overline{\mathbb{Q}}_\ell$-sheaf $\mathscr{L}_\psi$ of rank 1 on $\mathbb{A}^1_k$, which we call the Artin-Schreier sheaf. Let X be a scheme over k and let f be an element in the ring of global sections $\Gamma(X, \mathscr{O}_X)$ of the structure sheaf of X. Then f defines a k-morphism $X \to \mathbb{A}^1_k$ so that the induced k-algebra homomorphism $k[t] \to \Gamma(X, \mathscr{O}_X)$ maps t to f. We often denote this canonical morphism also by f, and denote by $\mathscr{L}_\psi(f)$ the inverse image of $\mathscr{L}_\psi$ under this morphism. Let $f_1, f_2 \in \Gamma(X, \mathscr{O}_X)$. We have

$$\mathscr{L}_\psi(f_1 - f_2) = \mathscr{L}_\psi(f_1) \otimes \mathscr{L}_\psi(f_2)^{-1}.$$

Moreover, we have $\mathscr{L}_\psi(f_1) \cong \mathscr{L}_\psi(f_2)$ if and only if $f_1 - f_2 = g^p - g$ for some $g \in \Gamma(X, \mathscr{O}_X)$. Here for the "only if" part, we need the assumption that $\psi : \mathbb{F}_p \to \overline{\mathbb{Q}}_l^*$ is nontrivial, which implies that ψ is injective. These facts will be used throughout this paper. (Confer [5] Sommes trig. 1.2-1.8.)

For any positive integer N prime to p, denote by $[N]$ the k-morphism

$$[N] : \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}, \quad t \mapsto t^N.$$

Let $I(k)$ be the set of positive integers N prime to p such that k contains a primitive N-th root of unity. For any $N \in I(k)$, $[N]$ is a finite Galois etale covering space. Let

$$\mu_N(k) = \{\mu \in k | \mu^N = 1\}.$$

Then $[N]$ defines a $\mu_N(k)$-torsor

$$1 \to \mu_N(k) \to \mathbb{G}_{m,k} \xrightarrow{[N]} \mathbb{G}_{m,k} \to 1.$$

When N runs over $I(k)$, we get an inverse system of extensions of $\mathbb{G}_{m,k}$ by $\mu_N(k)$. Passing to the limit, we get an extension of $\mathbb{G}_{m,k}$ by the profinite abelian group $\varprojlim_{N \in I(k)} \mu_N(k)$. Let

$$\rho : \varprojlim_{N \in I(k)} \mu_N(k) \to \mathrm{GL}(n, \overline{\mathbb{Q}}_\ell)$$

be a continuous representation. Pushing forward the above extension by ρ^{-1}, we get a lisse $\overline{\mathbb{Q}}_\ell$-sheaf $\mathscr{K}_\rho$ on $\mathbb{G}_{m,k}$ of rank n, which we call the Kummer sheaf associated to ρ. It is tamely ramified at 0 and at ∞. Conversely, when k is algebraically closed, any lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞ is obtained this way. This follows from [10] XIII 2.12. Let X be a scheme over k and let f be a unit in $\Gamma(X, \mathscr{O}_X)$. Then f defines a k-morphism $X \to \mathbb{G}_{m,k}$ so that the induced k-algebra homomorphism $k[t, 1/t] \to \Gamma(X, \mathscr{O}_X)$ maps t to f. We denote by $\mathscr{K}_\rho(f)$ the inverse image of $\mathscr{K}_\rho$ under this morphism.

Let $k((t))$ (resp. $k((1/t))$) be the field of formal Laurent series in the variable t (resp. $1/t$), and let $\eta_0 = \operatorname{Spec} k((t))$ (resp. $\eta_\infty = \operatorname{Spec} k((1/t))$). We have canonical morphisms $\eta_0 \to \mathbb{G}_m$ and $\eta_\infty \to \mathbb{G}_m$ defined by the inclusions $k[t, 1/t] \hookrightarrow k((t))$ and $k[t, 1/t] \hookrightarrow k((1/t))$, respectively. Objects on $\mathbb{G}_{m,k}$ can be restricted to η_0 and to η_∞ through these morphisms. By [18] 2.2.2.1, any tamely ramified $\overline{\mathbb{Q}}_\ell$-sheaf on η_0 (resp. η_∞) is the restriction of a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞.

For any positive integer r prime to p, denote also by $[r] : \eta_0 \to \eta_0$ the morphism induced by the k-algebra homomorphism

$$k((t)) \to k((t)), \quad t \mapsto t^r.$$

Any formal Laurent series

$$\alpha(t) = \frac{a_{-s}}{t^s} + \frac{a_{-(s-1)}}{t^{s-1}} + \cdots + \frac{a_{-1}}{t} + a_0 + a_1 t + \cdots$$

in $k((t))$ can be regarded as a global section of the structure sheaf of η_0. In this paper, we calculate the local Fourier transformation

$$\mathfrak{F}^{(0,\infty')}\big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big)\big)$$

for any Laurent series $\alpha(t)$ and any tamely ramified $\overline{\mathbb{Q}}_\ell$-sheaf $\mathscr{K}$ on η_0. Write

$$\alpha(t) = \alpha_1(t) + \alpha_2(t)$$

such that $\alpha_1(t)$ is the polar part of $\alpha(t)$ and $\alpha_2(t)$ is a formal power series of t. Then we have

$$[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big) \cong [r]_*\big(\mathscr{L}_\psi(\alpha_1(t)) \otimes \big(\mathscr{L}_\psi(\alpha_2(t)) \otimes \mathscr{K}\big)\big)$$

and $\mathscr{L}_\psi(\alpha_2(t)) \otimes \mathscr{K}$ is again tamely ramified. So to calculate $\mathscr{F}^{(0,\infty')}\big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big)\big)$ for all Laurent series $\alpha(t)$ and all tamely ramified sheaf $\mathscr{K}$, we may assume $\alpha(t)$ is a polynomial of $1/t$.

Similarly, let $[r] : \eta_\infty \to \eta_\infty$ be the morphism induced by the k-algebra homomorphism

$$k((1/t)) \to k((1/t)), \quad t \mapsto t^r.$$

Regard a formal Laurent series

$$\alpha(1/t) = a_{-s}t^s + a_{-(s-1)}t^{s-1} + \cdots + a_{-1}t + a_0 + \frac{a_1}{t} + \cdots$$

in $k((1/t))$ as a global section of the structure sheaf of η_∞. We also calculate the local Fourier transformations

$$\mathfrak{F}^{(\infty,0')}\big([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}\big)\big) \text{ and } \mathscr{F}^{(\infty,\infty')}\big([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}\big)\big)$$

for any Laurent series $\alpha(1/t)$ and any tamely ramified $\overline{\mathbb{Q}}_\ell$-sheaf $\mathscr{K}$ on η_∞. We refer the reader to [18] for the definitions and properties of local Fourier transformations.

Before presenting the main theorems of this paper, let's recall some facts from the classical analysis. Consider a smooth concave function $f(t)$ with $\frac{\mathrm{d}^2}{\mathrm{d}t^2}(f(t)) < 0$. The Legendre transformation of $f(t)$ is defined as follows: First consider the function

$$G(t, t') = f(t) + tt'.$$

Require the partial derivative $\frac{\partial G}{\partial t}$ vanishes:

$$\frac{\mathrm{d}}{\mathrm{d}t}(f(t)) + t' = 0.$$

Solve this equation for t as a function of t' and write the solution as $t = \mu(t')$. Note that for each fixed t', the function $G(t, t')$ reaches its greatest value at $t = \mu(t')$. The Legendre transformation L_f of f is defined to be

$$L_f(t') = G(\mu(t'), t') = f(\mu(t')) + \mu(t')t'.$$

(Classically the Legendre transformation is defined for convex functions $f(t)$ and $G(t, t')$ is taken to be $-f(t) + tt'$. Confer [2] Chap. 3, §14.) According to Laplace, (confer [4] Chapter 5, especially §17 and §20), under suitable conditions on $\phi(t, t')$, the major contribution to the value of the integral

$$\int_{-\infty}^{\infty} \phi(t, t')e^{f(t)+tt'}\mathrm{d}t$$

for large real t' comes from neighborhoods of those points where $G(t, t') = f(t)+tt'$ attains its greatest values, that is, those points on the curve $t = \mu(t')$. Moreover, the dominant term in the asymptotic

expansion of $\int_{-\infty}^{\infty}\phi(t,t')e^{f(t)+tt'}\mathrm{d}t$ is

$$\phi(\mu(t'),t')e^{L_f(t')}\sqrt{\frac{-\pi}{2\frac{\mathrm{d}^2 f}{\mathrm{d}t^2}(\mu(t'))}}.$$

Go back to the characteristic p case, and let

$$f(t)=\frac{a_{-s}}{(\sqrt[r]{t})^s}+\frac{a_{s-1}}{(\sqrt[r]{t})^{s-1}}+\cdots+\frac{a_{-1}}{\sqrt[r]{t}}+a_0+\cdots$$

be a Laurent series in the variable $\sqrt[r]{t}$ with $a_{-s}\neq 0$. Similar to the above discussion, we define the Legendre transformation $L_f(t')$ of $f(t)$ by the system of equations

$$\begin{cases} f(t)+tt'=L_f(t'), \\ \frac{\mathrm{d}}{\mathrm{d}t}(f(t))+t'=0. \end{cases}$$

Assume k is algebraically closed. Then $L_f(t')$ is a Laurent series in the variable $\frac{1}{\sqrt[r+s]{t'}}$ of the form

$$L_f(t')=b_{-s}(\sqrt[r+s]{t'})^s+b_{-(s-1)}(\sqrt[r+s]{t'})^{s-1}+\cdots+b_0+\frac{b_1}{\sqrt[r+s]{t'}}+\cdots$$

with $b_{-s}\neq 0$. Set

$$\alpha(t)=\sum_{i=-s}^{\infty}a_i t^i,\ \beta(1/t')=\sum_{i=-s}^{\infty}b_i(1/t')^i,$$

and let $\mathscr{L}(f)=[r]_*\mathscr{L}_\psi(\alpha(t))$ and $\mathscr{L}(L_f)=[r+s]_*\mathscr{L}_\psi(\beta(1/t'))$. Probably motivated by the asymptotic expansion discussed in the previous paragraph, Laumon and Malgrange conjecture that when $p\gg r,s$, we have

$$\mathfrak{F}^{(0,\infty')}(\mathscr{L}(f))\cong\mathscr{L}(L_f).$$

They also made similar conjectures for $\mathfrak{F}^{(\infty,0')}$ and $\mathfrak{F}^{(\infty,\infty')}$ ([18] 2.6.3). The main goal of this paper is to prove these conjectures (which need to be slightly corrected). Our main theorems are as follows:

Theorem 0.1. *Let $\gamma(t)=t^r$, let $\alpha(t)\in k[t,1/t]$, let*

$$\delta(t)=-\frac{1}{rt^{r-1}}\frac{\mathrm{d}}{dt}(\alpha(t)),$$
$$\beta(t')=\alpha(t')+\gamma(t')\delta(t'),$$

and let $\mathscr{K}$ be a rank 1 lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞. Denote by $\gamma,\delta:\mathbb{G}_{m,k}\to\mathbb{A}^1_k$ the k-morphisms defined by $\gamma(t)$ and $\delta(t')$, respectively.

(i) Suppose $\alpha(t)$ is of the form

$$\alpha(t)=\frac{a_{-s}}{t^s}+\frac{a_{-(s-1)}}{t^{s-1}}+\cdots+\frac{a_{-1}}{t}$$

with $a_{-s} \neq 0$, and suppose $r, s \geq 1$, $s < p$, and p is relatively prime to 2, r, s and $r+s$. We have

$$\mathfrak{F}^{(0,\infty')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big)\Big)|_{\eta_0}\Big) \cong \Big(\delta_*\Big(\mathscr{L}_\psi(\beta(t')) \otimes \mathscr{K} \otimes \mathscr{K}_{\chi_2}\Big(\frac{1}{2}s(r+s)a_{-s}t'^s\Big) \otimes G(\chi_2, \psi)\Big)\Big)|_{\eta_{\infty'}},$$

where $\mathscr{K}_{\chi_2}$ is the Kummer sheaf associated to the (unique) nontrivial character $\chi_2 : \mu_2(k) \to \overline{\mathbb{Q}}_\ell^*$ of order 2, and $G(\chi_2, \psi)$ is the unramified sheaf associated to the Gauss sum, that is, the $\overline{\mathbb{Q}}_\ell$-vector space of dimension 1 with continuous $\mathrm{Gal}(\bar{k}/k)$-action defined by

$$G(\chi_2, \psi) = H^1_c(\mathbb{G}_{m,k} \otimes_k \bar{k}, \mathscr{K}_{\chi_2} \otimes \mathscr{L}_\psi).$$

(ii) Suppose $\alpha(t)$ is of the form

$$\alpha(t) = a_s t^s + a_{s-1}t^{s-1} + \cdots + a_1 t$$

with $a_s \neq 0$, and suppose $r, s \geq 1$ and p is relatively prime to r and s. If $s \leq r$, then

$$\mathfrak{F}^{(\infty,\infty')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big)\Big)|_{\eta_\infty}\Big) = 0.$$

Suppose $s > r$ and suppose furthermore that $s < p$ and p is relatively prime to 2, r, s and $s-r$. We have

$$\mathfrak{F}^{(\infty,\infty')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big)\Big)|_{\eta_\infty}\Big) \cong \Big(\delta_*\Big(\mathscr{L}_\psi(\beta(t')) \otimes \mathscr{K} \otimes \mathscr{K}_{\chi_2}\Big(\frac{1}{2}s(s-r)a_s t'^s\Big) \otimes G(\chi_2, \psi)\Big)\Big)|_{\eta_{\infty'}}.$$

(iii) Suppose $\alpha(t)$ is of the form

$$\alpha(t) = a_s t^s + a_{s-1}t^{s-1} + \cdots + a_1 t$$

with $a_s \neq 0$, and suppose $r, s \geq 1$ and p is relatively prime to r and s. If $s \geq r$, then

$$\mathfrak{F}^{(\infty,0')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big)\Big)|_{\eta_\infty}\Big) = 0.$$

Suppose $s < r$ and suppose furthermore that $s < p$ and p is relatively prime to 2, r, s and $s-r$. We have

$$\mathfrak{F}^{(\infty,0')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big)\Big)|_{\eta_\infty}\Big) \cong \Big(\delta_*\Big(\mathscr{L}_\psi(\beta(t')) \otimes \mathscr{K} \otimes \mathscr{K}_{\chi_2}\Big(\frac{1}{2}s(s-r)a_s t'^s\Big) \otimes G(\chi_2, \psi)\Big)\Big)|_{\eta_{0'}}.$$

In the case where k is algebraically closed, we have the following theorems:

Theorem 0.2. *Suppose k is algebraically closed, $r, s \geq 1$, $s < p$, and p is relatively prime to 2, r, s and $r+s$. Let*

$$\alpha(t) = \sum_{i=-s}^{\infty} \frac{a_i}{t^i}$$

be a formal Laurent series in $k((1/t))$ with $a_{-s} \neq 0$. Consider the system of equations

$$\begin{cases} \alpha(t) + t^r t'^{r+s} = \beta(1/t'), \\ \frac{\mathrm{d}}{\mathrm{d}t}(\alpha(t)) + r t^{r-1} t'^{r+s} = 0. \end{cases}$$

Using the second equation, we find an expression of t in terms of t'. We then substitute this expression into the first equation to get $\beta(1/t')$, which is a formal Laurent series in $k((1/t'))$ of the form

$$\beta(1/t') = \sum_{i=-s}^{\infty} b_i (1/t')^i$$

with $b_{-s} \neq 0$. For any lisse $\overline{\mathbb{Q}}_\ell$-sheaf $\mathscr{K}$ on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞, we have

$$\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}|_{\eta_0}\big)\Big) \cong [r+s]_*\big(\mathscr{L}_\psi(\beta(1/t')) \otimes (\mathrm{inv}^*\mathscr{K})|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big),$$

where $\mathrm{inv} : \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$ is the morphism $t' \mapsto \frac{1}{t'}$, and $\mathscr{K}_{\chi_2}$ is the Kummer sheaf associated to the (unique) nontrivial character $\chi_2 : \mu_2(k) \to \overline{\mathbb{Q}}_\ell^*$ of order 2.

Theorem 0.3. *Suppose k is algebraically closed, $r, s \geq 1$ and p is relatively prime to r and s. Let*

$$\alpha(1/t) = \sum_{i=-s}^{\infty} a_i (1/t)^i$$

be a formal Laurent series in $k((1/t))$ with $a_{-s} \neq 0$, and let $\mathscr{K}$ be a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞. If $s \leq r$, then

$$\mathfrak{F}^{(\infty,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}|_{\eta_\infty}\big)\Big) = 0.$$

Suppose $s > r$ and suppose furthermore that $s < p$ and p is relatively prime to 2, r, s and $s - r$. Define $\beta(1/t')$ by the system of equations

$$\begin{cases} \alpha(1/t) + t^r t'^{s-r} = \beta(1/t'), \\ \frac{\mathrm{d}}{\mathrm{d}t}(\alpha(1/t)) + r t^{r-1} t'^{s-r} = 0. \end{cases}$$

It is a formal Laurent series in $k((1/t'))$ of the form

$$\beta(1/t') = \sum_{i=-s}^{\infty} b_i (1/t')^i$$

with $b_{-s} \neq 0$. We have

$$\mathfrak{F}^{(\infty,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}|_{\eta_\infty}\big)\Big) \cong [s-r]_*\big(\mathscr{L}_\psi(\beta(1/t')) \otimes \mathscr{K}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big),$$

where $\mathscr{K}_{\chi_2}$ is the Kummer sheaf associated to the (unique) nontrivial character $\chi_2 : \mu_2(k) \to \overline{\mathbb{Q}}_\ell^*$ of order 2.

Theorem 0.4. *Suppose k is algebraically closed, $r, s \geq 1$ and p is relatively prime to r and s. Let*

$$\alpha(1/t) = \sum_{i=-s}^{\infty} a_i (1/t)^i$$

be a formal Laurent series in $k((1/t))$ with $a_{-s} \neq 0$, and let $\mathscr{K}$ be a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞. If $s \geq r$, then

$$\mathfrak{F}^{(\infty,0')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}|_{\eta_\infty}\big)\Big) = 0.$$

Suppose $s < r$ and suppose furthermore that $s < p$ and p is relatively prime to 2, r, s and $r - s$. Define $\beta(t')$ by the system of equations

$$\begin{cases} \alpha(1/t) + t^r t'^{r-s} = \beta(t'), \\ \frac{\mathrm{d}}{\mathrm{d}t}(\alpha(1/t)) + r t^{r-1} t'^{r-s} = 0. \end{cases}$$

It is a formal Laurent series in $k((t'))$ of the form

$$\beta(t') = \sum_{i=-s}^{\infty} b_i t'^i$$

with $b_{-s} \neq 0$. We have

$$\mathfrak{F}^{(\infty,0')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}|_{\eta_\infty}\big)\Big) \cong [r-s]_*\big(\mathscr{L}_\psi(\beta(t')) \otimes (\mathrm{inv}^*\mathscr{K})|_{\eta_{0'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{0'}}\big),$$

where $\mathrm{inv} : \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$ *is the morphism $t' \mapsto \frac{1}{t'}$, and $\mathscr{K}_{\chi_2}$ is the Kummer sheaf associated to the (unique) nontrivial character $\chi_2 : \mu_2(k) \to \overline{\mathbb{Q}}_\ell^*$ of order 2.*

When $\mathscr{K}$ is trivial, Theorems 0.2-0.4 are conjectured by Laumon and Malgrange except that the term $[s]^*\mathscr{K}_{\chi_2}$ is missing in their conjecture. Note that in these theorems, the system of equations relating α and β is equivalent to the system of equations defining the Legendre transformation. For example, in Theorem 0.2, $\beta(1/\sqrt[r+s]{t'})$ is the Legendre transformation of $\alpha(\sqrt[r]{t})$.

Note that we take $\mathscr{K}$ to be of rank 1 if k is not assumed to be algebraically closed, whereas $\mathscr{K}$ has arbitrary rank if k is algebraically closed. This is for the technical reason that when k is algebraically closed, we know the tame fundamental group of $\mathbb{G}_{m,k}$ is isomorphic to $\varprojlim_{(N,p)=1} \mu_N(k)$ so that we have a structure theorem (Lemma 2.5) for lisse $\overline{\mathbb{Q}}_\ell$-sheaves on $\mathbb{G}_{m,k}$ tamely ramified at 0 and ∞. This is not available when k is not algebraically closed.

In [19], Ramero constructs the Fourier transformations for a class of étale analytic sheaves on the analytification of the affine line over a field of characteristic 0, complete with respect to a non-Archimedean metric. In order for the Fourier transformations to be constructible sheaves, the analytic

sheaves are assumed to have meromorphic ramification ([19] 8.3.6). Ramero asks whether the Fourier transformation of an analytic sheaf with meromorphic ramification still has meromorphic ramification, and this problem can be translated into a problem concerning the local Fourier transformation of a meromorphic representation ([19] 8.7). An analytic sheaf with meromorphic ramification on the germ of a punctured disc at 0 is nothing but an analytic analogue of the $\overline{\mathbb{Q}}_\ell$-sheaf $[r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}|_{\eta_0}\big)$ studied in this paper (confer [19] §5 and 8.1.1). The method used in this paper can also be used to calculate the local Fourier transformations of analytic sheaves with meromorphic ramification. The resulting formulas are the same as those in Theorems 0.2-0.4. In particular, we can give a positive answer to Ramero's question asking whether the Fourier transformation of an analytic sheaf with meromorphic ramification still has meromorphic ramification. In fact, it was for the purpose of solving Ramero's problem that I started working on the Laumon-Malgrange conjecture.

About the same time when the main results of this paper was obtained, analogous formulas for the local Fourier transformations of D-modules were proved independently by J. Fang [8] and C. Sabbah [20]. By the refined Turrittin-Levelt Theorem ([20] Corollary 3.3), when k is an algebraically closed field of characteristic 0, any finite dimensional $k((t))$-vector space M with a connection is a direct sum of the D-module analogues of the $\overline{\mathbb{Q}}_\ell$-sheaves $[r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}|_{\eta_0}\big)$. So to calculate the local Fourier transformation of M, it suffices to calculate the local Fourier transformations of the D-module analogues of $[r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}|_{\eta_0}\big)$. Of course, the Turrittin-Levelt Theorem doesn't hold for $\overline{\mathbb{Q}}_\ell$-sheaves on η_0 in the characteristic p case. But Deligne ([6]) suggested to me the following analogue of the Turrittin-Levelt Theorem in characteristic p.

Proposition 0.5. *Suppose k is algebraically closed. Let $\rho : I \to \mathrm{GL}(V)$ be an irreducible $\overline{\mathbb{Q}}_\ell$-representation of $I = \mathrm{Gal}\Big(\overline{k((t))}/k((t))\Big)$, where $\overline{k((t))}$ is a separable closure of $k((t))$. Suppose the following conditions hold:*

(a) $\rho(P^p[P,P]) = 1$, where $P = \mathrm{Gal}\left(\overline{k((t))}/\bigcup_{(N,p)=1} k(\sqrt[N]{t})\right)$ is the wild inertia subgroup.

(b) $\rho(I)$ is finite.

(c) Let s be the Swan conductor of ρ. We have $s < p$.

Then there exist a character $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$ of finite order, a Laurent series $\alpha(t) \in k((t))$ of the form

$$\alpha(t) = \frac{a_{-s}}{t^s} + \frac{a_{-(s-1)}}{t^{s-1}} + \cdots + \frac{a_{-1}}{t}$$

with $a_{-s} \neq 0$, and a positive integer r prime to p such that V is isomorphic to the representation

corresponding to the lisse $\overline{\mathbb{Q}}_\ell$*-sheaf* $[r]_*(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi)$ *on* $\eta_0 = \operatorname{Spec} k((t))$.

In the above proposition, condition (a) is necessary for the representation V to be isomorphic to the representation corresponding to a sheaf of the form $[r]_*(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi)$ on $\eta_0 = \operatorname{Spec} k((t))$. Condition (b) holds for most interesting representations. For example, if we know ρ is irreducible and is quasi-unipotent, that is, there exists a closed subgroup I_1 of I of finite index such that $\rho(\sigma)$ is unipotent for any $\sigma \in I_1$, then (b) holds. We know representations coming from arithmetic and geometry are often quasi-unipotent. See for example, the appendix of [22], and [11] I 1.3. Condition (c) should not be essential. But we need it in our proof for some algebraic argument to work.

In the following, $\mathbb{Z}$-schemes and $\mathbb{Z}$-morphisms are assumed to be separated and of finite type. Taking fibers at a prime p of $\mathbb{Z}$-schemes and $\mathbb{Z}$-morphisms is indicated by putting the subscript p. In [16] pages 297-301, in order to have a general framework to study the variation with p of exponential sums on schemes over $\mathbb{Z}$, Katz proposes to study the family

$$\{(Y_p, \mathscr{G}_p) | p \text{ is prime}\},$$

where Y is a scheme over $\mathbb{Z}$, and for each p prime to ℓ, $\mathscr{G}_p$ is a $\overline{\mathbb{Q}}_\ell$-sheaf on the fiber Y_p so that there exist a scheme X over $\mathbb{Z}$, an object K in the category $D_c^b(X, \overline{\mathbb{Q}}_\ell)$ of complexes of $\overline{\mathbb{Q}}_\ell$-sheaves on X constructed in [7] 1.1, and two $\mathbb{Z}$-morphisms $f : X \to Y$ and $g : X \to \mathbb{A}^1_{\mathbb{Z}}$ with the property

$$\mathscr{G}_p \cong R^i f_{p!}(K|_{X_p} \otimes g_p^* \mathscr{L}_{\psi_p})$$

for some i and all p prime to ℓ, where $\mathscr{L}_{\psi_p}$ is the Artin-Schreier sheaf defined by the standard additive character

$$\psi_p : \mathbb{F}_p \to \overline{\mathbb{Q}}_l^*, \quad \psi_p(t) = e^{\frac{2\pi i t}{p}}.$$

Let

$$Y \overset{\mathrm{pr}_1}{\leftarrow} Y \times \mathbb{A}^1_{\mathbb{Z}} \overset{\mathrm{pr}_2}{\to} \mathbb{A}^1_{\mathbb{Z}}$$

be the projections, let $h : X \to Y \times \mathbb{A}^1_{\mathbb{Z}}$ be the morphism defined by

$$\mathrm{pr}_1 \circ h = f, \ \mathrm{pr}_2 \circ h = g,$$

and let $L = Rh_! K$. Then we have

$$\mathscr{G}_p \cong R^i \mathrm{pr}_{1p!}(L|_{(Y \times \mathbb{A}^1_{\mathbb{Z}})_p} \otimes \mathrm{pr}_{2p}^* \mathscr{L}_{\psi_p}).$$

So in the definition of the family $\{(Y_p, \mathscr{G}_p)\}$, we may require $X = Y \times \mathbb{A}^1_{\mathbb{Z}}$, $f = \mathrm{pr}_1$ and $g = \mathrm{pr}_2$.

Conjecture 0.6. *With the above notations, suppose Y is an algebraic curve over $\mathbb{Z}$. Let $\overline{\mathbb{F}}_p$ be an algebraic closure of $\mathbb{F}_p$. Then for sufficiently large p, the inverse image of $\mathscr{G}_p$ by any dominant morphism*

$$\mathrm{Spec}\,\overline{\mathbb{F}}_p((t)) \to Y_p$$

can be written as a direct sum of objects of the form

$$[r]_*(\mathscr{L}_{\psi_p}(\alpha(t)) \otimes \mathscr{K}),$$

where r is relatively prime to p, $\alpha(t) \in \overline{\mathbb{F}}_p((t))$, and $\mathscr{K}$ is a tame $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathrm{Spec}\,\overline{\mathbb{F}}_p((t))$.

We expect the method used in this paper is helpful to prove the above conjecture. In any case, Theorems 0.1-0.4 can be used to calculate the local monodromy of sheaves arising from the study of exponential sums. Indeed, as a direct application of our theorems, we calculate the local monodromy of hypergeometric $\overline{\mathbb{Q}}_\ell$-sheaves introduced by Katz ([15]).

Let

$$\lambda_1, \ldots, \lambda_n, \rho_1, \ldots, \rho_m : \mathbb{F}_q^* \to \overline{\mathbb{Q}}_\ell^*$$

be multiplicative characters. For any extension $\mathbb{F}_{q^k}$ of $\mathbb{F}_q$ of degree k, and any $t \neq 0$ in $\mathbb{F}_{q^k}$, we define the hypergeometric sum by

$$\begin{aligned}
&\mathrm{Hyp}(\psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)(\mathbb{F}_{q^k}, t) \\
= &\sum_{x_1,\ldots,x_n,y_1,\ldots,y_m \in \mathbb{F}_{q^k}, x_1\cdots x_n = t y_1 \cdots y_m} \psi\Big(\mathrm{Tr}_{\mathbb{F}_{q^k}/\mathbb{F}_p}(\sum_{i=1}^n x_i - \sum_{j=1}^m y_j)\Big) \\
&\qquad \times \prod_{i=1}^n \lambda_i\Big(\mathrm{N}_{\mathbb{F}_{q^k}/\mathbb{F}_q}(x_i)\Big) \prod_{j=1}^m \rho_j\Big(\mathrm{N}_{\mathbb{F}_{q^k}/\mathbb{F}_q}\Big(\frac{1}{y_j}\Big)\Big).
\end{aligned}$$

In the case where $m = 0$, we get the Kloosterman sum

$$\mathrm{Kl}_n(\psi; \lambda_1, \ldots, \lambda_n)(\mathbb{F}_{q^k}, t) = \sum_{x_1,\ldots,x_n \in \mathbb{F}_{q^k}, x_1\cdots x_n = t} \psi\Big(\mathrm{Tr}_{\mathbb{F}_{q^k}/\mathbb{F}_p}(\sum_{i=1}^n x_i)\Big) \prod_{i=1}^n \lambda_i\Big(\mathrm{N}_{\mathbb{F}_{q^k}/\mathbb{F}_q}(x_i)\Big).$$

In [15] 8.2, Katz constructs an object $\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)$ in the triangulated category $D_c^b(\mathbb{G}_{m,\mathbb{F}_q}, \overline{\mathbb{Q}}_\ell)$ so that for any $\mathbb{F}_{q^k}$-rational point t of $\mathbb{G}_{m,\mathbb{F}_q}$, we have

$$\mathrm{Tr}(F_t, (\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m))_{\bar{t}}) = (-1)^{n+m}\mathrm{Hyp}(\psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)(\mathbb{F}_{q^k}, t),$$

where F_t is the geometric Frobenius element at t. In fact, Katz constructs $\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)$ on $\mathbb{G}_{m,k}$ for any field k of characteristic p and any characters

$$\lambda_1, \ldots, \lambda_n, \rho_1, \ldots, \rho_m : \varprojlim_{N \in I(k)} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*.$$

Assume k is algebraically closed. Fix an element

$$\zeta = (\zeta_N) \in \varprojlim_{(N,p)=1} \mu_N(k)$$

so that ζ_N is a primitive N-th root of unity for each N prime to p. Note that ζ is a topological generator of $\varprojlim_{(N,p)=1} \mu_N(k)$. Let $U(n)$ be an n-dimensional vector space $\overline{\mathbb{Q}}_\ell^n$ on which ζ acts through the unipotent $(n \times n)$-matrix with a single Jordan block

$$\begin{pmatrix} 1 & 1 & & \\ & 1 & \ddots & \\ & & \ddots & 1 \\ & & & 1 \end{pmatrix}.$$

This action can be extended to a continuous action of $\varprojlim_{(N,p)=1} \mu_N(k)$ on $U(n)$. Denote also by $U(n)$ the corresponding lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞. For sufficiently large p, the following proposition is more precise than [15] 8.4.2 (6) and (7).

Proposition 0.7. *Let k be an algebraically closed field of characteristic p, and let*

$$\lambda_1, \ldots, \lambda_n, \rho_1, \ldots, \rho_m : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$$

be characters. For any character $\lambda : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$, let $\mathrm{mult}_0(\lambda)$ be the number of times of λ occurring in $\lambda_1, \ldots, \lambda_n$, and let $\mathrm{mult}_\infty(\lambda)$ be the number of times of λ occurring in $\rho_1, \ldots, \rho_m$. Suppose the family of characters $\lambda_1, \ldots, \lambda_n$ is disjoint from the family of characters $\rho_1, \ldots, \rho_m$, and suppose p is relatively prime to $2, 3, \ldots, \max(m, n)$.

(i) If $n > m$, then $\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)[-1]$ is an irreducible lisse sheaf of rank n on $\mathbb{G}_{m,k}$, and

$$\begin{aligned} &\big(\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)[-1]\big)|_{\eta_0} \cong \bigoplus_\lambda \big(\mathscr{K}_\lambda \otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0}, \\ &\big(\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)[-1]\big)|_{\eta_\infty} \\ \cong\ & [n-m]_*\big(\mathscr{L}_\psi((n-m)t) \otimes \mathscr{K}_{\lambda_1 \cdots \lambda_n \rho_1^{-1} \cdots \rho_m^{-1}} \otimes \mathscr{K}_{\chi_2{}^{n+m-1}}\big)|_{\eta_\infty} \bigoplus \bigoplus_\rho \big(\mathscr{K}_\rho \otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty}. \end{aligned}$$

(ii) If $n < m$, then $\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)[-1]$ is an irreducible lisse sheaf of rank m on

$\mathbb{G}_{m,k}$, *and*

$$\begin{aligned}
&\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty} \cong \bigoplus_\rho \big(\mathscr{K}_\rho \otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty},\\
&\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_0}\\
\cong\ & [m-n]_*\left(\mathscr{L}_\psi\left(-\frac{m-n}{t}\right)\otimes \mathscr{K}_{\lambda_1^{-1}\cdots\lambda_n^{-1}\rho_1\cdots\rho_m}\otimes \mathscr{K}_{\chi_2{}^{n+m-1}}\right)|_{\eta_0}\bigoplus\bigoplus_\lambda \big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0}.
\end{aligned}$$

(iii) If $n=m$, *then* $\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m)[-1]$ *is an irreducible lisse sheaf of rank* n *when restricted to* $\mathbb{G}_{m,k}-\{1\}$, *and we have*

$$\begin{aligned}
\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_0} &\cong \bigoplus_\lambda \big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0},\\
\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty} &\cong \bigoplus_\rho \big(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty}.
\end{aligned}$$

Let $k((t-1))$ *be the field of Laurent power series in the variable* $t-1$, *let* $\eta_1=\mathrm{Spec}\,k((t-1))$, *and let*

$$\mathrm{tran}:\mathbb{A}^1_k\to\mathbb{A}^1_k,\quad x\mapsto x-1$$

be the translation by 1. *We have*

$$(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_n)[-1])_{\bar 1}\cong(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_n)[-1])_{\bar\eta_1}^{\mathrm{Gal}(\bar\eta_1/\eta_1)},$$

and we have a short exact sequence

$$\begin{aligned}
0 &\to (\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_n)[-1])_{\bar\eta_1}^{\mathrm{Gal}(\bar\eta_1/\eta_1)}\\
&\to (\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_n)[-1])_{\bar\eta_1}\\
&\to (\mathrm{tran}^*\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_n^{-1}\rho_1\cdots\rho_n})|_{\bar\eta_1}\to 0.
\end{aligned}$$

Taking $m=0$, we get the following.

Proposition 0.8. *Suppose* k *is algebraically closed and* $p=\mathrm{char}\,k$ *is relatively prime to* $2,3,\dots,n$. *The Kloosterman sheaf*

$$\mathrm{Kl}_n(\psi;\lambda_1,\dots,\lambda_n)=\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\emptyset)[-1]$$

is an irreducible lisse sheaf of rank n *on* $\mathbb{G}_{m,k}$ *and*

$$\begin{aligned}
\mathrm{Kl}_n(\psi;\lambda_1,\dots,\lambda_n)|_{\eta_0} &\cong \bigoplus_\lambda\big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0},\\
\mathrm{Kl}_n(\psi;\lambda_1,\dots,\lambda_n)|_{\eta_\infty} &\cong [n]_*(\mathscr{L}_\psi(nt)\otimes\mathscr{K}_{\lambda_1\cdots\lambda_n}\otimes\mathscr{K}_{\chi_2{}^{n-1}})|_{\eta_\infty}.
\end{aligned}$$

When $\lambda_1, \ldots, \lambda_n$ are trivial, the above assertion about $\mathrm{Kl}_n(\psi; \lambda_1, \ldots, \lambda_n)|_{\eta_\infty}$ is Lemma 1.6 in [9]. However in [9], we only assume p is relatively prime to n. The result in [9] is based on the work of Katz ([17] 10.1 and 5.6.2). Katz deduces his results from an identity of Hasse-Davenport. In [15] 8.4.2 (8), Katz shows that Proposition 0.7 (iii) holds without the assumption that p is relatively prime to $1, \ldots, n$. It is an interesting question to determine explicitly the local monodromy at 0 and at ∞ for $\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)$ without assuming that p is relatively prime to $2, 3, \ldots, \max(n, m)$. In fact, I don't know $\mathrm{Kl}_2(\psi; 1, 1)|_{\eta_\infty}$ explicitly if $p = 2$.

The paper is organized as follows. In §1, we calculate a direct factor of the restriction of the global Fourier transformation at $\eta_{\infty'}$. In §2, we use a group theoretical argument to deduce Theorems 0.1-0.4 from results in §1, and we prove Proposition 0.5 at the end of this section. We prove Proposition 0.7 in §3.

Shortly after the first version of the paper was finished, Abbes and Saito [1] were able to calculate the local Fourier transformations for a class of monomial Galois representations of Artin–Schreier-Witt type. Their results are more general than ours, and they used a blowing-up technique and the ramification theory of Kato. The method used in this paper is more direct and global. In the first version of this paper, we work over an algebraically closed field k and prove Theorems 0.2-0.4. Theorems 0.1 is inspired by [1] and is suggested by the referee.

1 Key lemmas

The main result of this section is Lemma 1.5, which calculates a direct factor of

$$\Big(R^1\pi_{2!}(\mathscr{L}_\psi(f(x,t')) \otimes \pi_1^*\mathscr{K})\Big)|_{\eta'_\infty},$$

where $\pi_1, \pi_2 : \mathbb{G}_{m,k} \times_k \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$ are the projections, $f(x,t') \in k[x, 1/x, t', 1/t']$ satisfies certain conditions and $\mathscr{K}$ is a rank 1 lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ such that $\mathscr{K}|_1 = \overline{\mathbb{Q}}_\ell$. Here we consider the rational point 1 of $\mathbb{G}_{m,k}$ as a closed subscheme of $\mathbb{G}_{m,k}$.

Lemma 1.1. *Let R be a commutative ring and let $h(x) = \sum_{i=-m}^{n} c_i x^i$ be a Laurent polynomial with $c_i \in R$ and $m, n \geq 0$.*

(i) Suppose $m, n \geq 1$ and c_n and c_{-m} are units in R. Then the R-morphism

$$H : \mathbb{G}_{m,R} \to \mathbb{A}^1_R, \quad x \mapsto h(x)$$

corresponding to the R-algebra homomorphism

$$R[x] \to R[x, 1/x], \quad x \mapsto h(x)$$

is a finite morphism, and it can be extended to an R-morphism

$$\hat{H} : \mathbb{P}^1_R \to \mathbb{P}^1_R$$

which in terms of homogenous coordinates can be described by

$$[x_0 : x_1] \mapsto [x_0^n x_1^m : \sum_{i=-m}^{n} c_i x_0^{n-i} x_1^{m+i}].$$

The canonical diagram

$$\begin{array}{ccc} \mathbb{G}_{m,R} & \hookrightarrow & \mathbb{P}^1_R \\ {\scriptstyle H}\downarrow & & \downarrow{\scriptstyle \hat{H}} \\ \mathbb{A}^1_R & \hookrightarrow & \mathbb{P}^1_R \end{array}$$

is Cartesian.

(ii) Suppose $m = 0$, $n \geq 1$ and c_n is a unit in R. Then the R-morphism

$$H : \mathbb{A}^1_R \to \mathbb{A}^1_R, \quad x \mapsto h(x)$$

is a finite morphism, and it can be extended to an R-morphism

$$\hat{H} : \mathbb{P}^1_R \to \mathbb{P}^1_R$$

which in terms of homogenous coordinates can be described by

$$[x_0 : x_1] \mapsto [x_0^n : \sum_{i=0}^{n} c_i x_0^{n-i} x_1^i].$$

The canonical diagram

$$\begin{array}{ccc} \mathbb{A}^1_R & \hookrightarrow & \mathbb{P}^1_R \\ {\scriptstyle H}\downarrow & & \downarrow{\scriptstyle \hat{H}} \\ \mathbb{A}^1_R & \hookrightarrow & \mathbb{P}^1_R \end{array}$$

is Cartesian.

Proof. We sketch a proof of (i). Consider the R-algebra homomorphism of graded rings

$$\begin{aligned} &R[x_0, x_1] \to R[x_0, x_1], \\ &x_0 \mapsto x_0^n x_1^m, \quad x_1 \mapsto \sum_{i=-m}^{n} c_i x_0^{n-i} x_1^{m+i}. \end{aligned}$$

This homomorphism does not preserve degrees, but maps a homogenous polynomial of degree d to a homogenous polynomial of degree $(m+n)d$. However by [12] 2.4.7 (i) and 2.8.2, it still defines an

R-morphism $\hat{H} : \mathbb{P}^1_R \to \mathbb{P}^1_R$. Here we need the fact that c_n and c_{-m} are units in order for the domain of definition of $\hat{H}$ to be the whole projective line $\mathbb{P}^1_R$. One can verify that $\hat{H}^{-1}(\mathbb{A}^1_R) = \mathbb{G}_{m,R}$. This implies the diagram in the lemma is Cartesian. So H is a proper morphism. By definition, H is an affine morphism. So H is a finite morphism by [13] 4.4.2. □

Lemma 1.2. *Let $g(x, z') \in k[x, 1/x, z']$, and let $a_0(x) \in k[x, 1/x]$ be the constant term of $g(x, z')$ considered as a polynomial of z'. Write $g(x, z')$ as*

$$g(x, z') = \sum_{i=-m}^{n} c_i(z')x^i,$$

where $m, n \geq 0$ and $c_i(z') \in k[z']$. Let $\mathscr{K}$ be a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$, and let $\bar{k}$ be an algebraic closure of k.

(i) Suppose $m, n \geq 1$ and $c_n(0)c_{-m}(0) \neq 0$. Denote by G the k-morphism

$$G : \mathbb{G}_{m,k} \times_k \mathbb{A}^1_k \to \mathbb{A}^1_k \times_k \mathbb{A}^1_k, \quad (x, z') \to (g(x, z'), z').$$

Let

$$S = \{a \in \bar{k} \mid a = a_0(b) \text{ for some } b \in \bar{k} - \{0\} \text{ satisfying } \frac{\mathrm{d}a_0}{\mathrm{d}x}(b) = 0\}.$$

For any $x \in \bar{k} - S$, there exists a neighborhood of $(x, 0)$ in $\mathbb{A}^1_k \times_k \mathbb{A}^1_k$ in which $G_!\pi_1^*\mathscr{K}$ is lisse and $R^iG_!\pi_1^*\mathscr{K}$ $(i \geq 1)$ vanish, where $\pi_1 : \mathbb{G}_{m,k} \times \mathbb{A}^1_k \to \mathbb{G}_{m,k}$ is the projection.

(ii) Suppose $m = 0$, $n \geq 1$ and $c_n(0) \neq 0$. Denote by G the k-morphism

$$G : \mathbb{A}^1_k \times_k \mathbb{A}^1_k \to \mathbb{A}^1_k \times_k A^1_k, \quad (x, z') \to (g(x, z'), z').$$

Let

$$S = \{a \in \bar{k} | a = a_0(b) \text{ for some } b \in \bar{k} \text{ satisfying } \frac{\mathrm{d}a_0}{dx}(b) = 0\} \cup \{a_0(0)\}.$$

For any $x \in \bar{k} - S$, there exists a neighborhood of $(x, 0)$ in $\mathbb{A}^1_k \times_k \mathbb{A}^1_k$ in which $G_!p_1^*j_!\mathscr{K}$ is lisse and $R^iG_!p_1^*j_!\mathscr{K}$ $(i \geq 1)$ vanish, where $j : \mathbb{G}_{m,k} \hookrightarrow \mathbb{A}^1_k$ is the open immersion, and $p_1 : \mathbb{A}^1_k \times_k \mathbb{A}^1_k \to \mathbb{A}^1_k$ is the projection to the first factor.

Proof. The proofs of (i) and (ii) are similar. We give a proof (i). The Jacobian G is

$$\det \begin{pmatrix} \frac{\partial}{\partial x}(g(x, z')) & \frac{\partial}{\partial z'}(g(x, z')) \\ \frac{\partial}{\partial x}(z') & \frac{\partial}{\partial z'}(z') \end{pmatrix} \equiv \frac{\mathrm{d}a_0}{\mathrm{d}x} \mod z'.$$

For $z' = 0$, the above Jacobian vanishes if and only if $\frac{\mathrm{d}a_0}{\mathrm{d}x} = 0$. So $G : \mathbb{G}_{m,k} \times_k \mathbb{A}^1_k \to \mathbb{A}^1_k \times_k \mathbb{A}^1_k$ is étale at $(b, 0)$ if $\frac{\mathrm{d}a_0}{\mathrm{d}x}(b) \neq 0$. Hence G is étale at every preimage of $(x, 0)$ if $x \notin S$. Next we prove G is

a finite morphism over some neighborhood of 0 in $\mathbb{A}^1_k$. This will imply (i). Set $A = k[z']$. Then G is the morphism

$$G : \mathbb{G}_{m,A} \to \mathbb{A}^1_A$$

corresponding to the A-algebra homomorphism

$$A[x] \to A[x, 1/x], \quad x \mapsto g(x, z').$$

Let $R = A_{c_n c_{-m}}$ be the localization of A with respect to $c_n(z')c_{-m}(z') \in A$. Then $c_n(z')$ and $c_{-m}(z')$ are unit in R. By Lemma 1.1 (i), $G : \mathbb{G}_{m,R} \to \mathbb{A}^1_R$ is a finite morphism. Since $c_n(0)c_{-m}(0) \neq 0$, 0 is contained in the image of the open immersion

$$\operatorname{Spec} R = \operatorname{Spec} A_{c_n c_{-m}} \hookrightarrow \operatorname{Spec} A = \mathbb{A}^1_k.$$

Our assertion follows. □

From now on we assume the characteristic p of k is distinct from 2.

Lemma 1.3. *Let $\mathcal{O}^{\mathrm{h}}_{\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k, (1,0)}$ (resp. $\mathcal{O}^{\mathrm{h}}_{\mathbb{A}^1_k \times_k \mathbb{A}^1_k, (0,0)}$) be the henselization of $\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k$ (resp. $\mathbb{A}^1_k \times_k \mathbb{A}^1_k$) at the point $(1,0)$ (resp. $(0,0)$), let $(\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(1,0)}$ (resp. $(\mathbb{A}^1_k \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(0,0)}$) be their spectrum, let*

$$\begin{aligned} G : \mathbb{G}_{m,k} \times_k \mathbb{A}^1_k &\to \mathbb{A}^1_k \times_k \mathbb{A}^1_k, \\ (x, z') &\mapsto (g(x, z'), z') \end{aligned}$$

be the k-morphism defined by some $g(x, z') \in k[x, 1/x, z']$ with the property

$$\begin{aligned} &g(1, z') = 0, \\ &\frac{\partial g}{\partial x}(1, z') = 0, \\ &\frac{1}{2}\frac{\partial^2 g}{\partial x^2}(1, 0) \text{ is a nonzero square in } k, \end{aligned}$$

and let

$$\omega : (\mathbb{A}^1_k \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(0,0)} \to (\mathbb{A}^1_k \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(0,0)}$$

be the morphism induced on henselizations by the morphism

$$\mathbb{A}^1_k \times_k \mathbb{A}^1_k \to \mathbb{A}^1_k \times_k \mathbb{A}^1_k, \quad (x, z') \mapsto (x^2, z').$$

Then there exists an isomorphism

$$\phi : (\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(1,0)} \xrightarrow{\cong} (\mathbb{A}^1_k \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(0,0)}$$

such that the morphism on henselizations

$$G^{\mathrm{h}} : (\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(1,0)} \to (\mathbb{A}^1_k \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(0,0)}$$

induced by G coincides with $\omega \circ \phi$.

Proof. Since $g(1, z') = \frac{\partial g}{\partial x}(1, z') = 0$, we have $\frac{g(x,z')}{(x-1)^2} \in k[x, 1/x, z']$. Hence $\frac{g(x,z')}{(x-1)^2}$ lies in $\mathscr{O}^{\mathrm{h}}_{\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k,(1,0)}$ and its image in the residue field is $\frac{1}{2}\frac{\partial^2 g}{\partial x^2}(1, 0)$, which is a nonzero square in k. Since $\operatorname{char} k \neq 2$, this element in the residue field has two distinct square roots. By the Hensel lemma [14] 18.5.13, $\frac{g(x,z')}{(x-1)^2}$ has two distinct square roots in $\mathscr{O}^{\mathrm{h}}_{\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k,(1,0)}$. Let $\delta(x - 1, z')$ be one of the square roots. Its image in the residue field is nonzero, and we have

$$g(x, z') = ((x - 1)\delta(x - 1, z'))^2.$$

Consider the morphism

$$(\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(1,0)} \to \mathbb{A}^1_k \times_k \mathbb{A}^1_k$$

corresponding to the k-algebra homomorphism

$$\begin{aligned} &k[x, z'] \to \mathscr{O}^{\mathrm{h}}_{\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k,(1,0)}, \\ &x \mapsto (x - 1)\delta(x - 1, z'), \quad z' \mapsto z'. \end{aligned}$$

It maps the Zariski closed point of $(\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(1,0)}$ to $(0, 0)$. Denote the induced morphism on henselizations by

$$\phi : (\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(1,0)} \to (\mathbb{A}^1_k \times_k \mathbb{A}^1_k)^{\mathrm{h}}_{(0,0)}.$$

Then $G^{\mathrm{h}} = \omega \circ \phi$. Let's prove ϕ is an isomorphism. By [14] 17.6.3, it suffices to the homomorphism induced by ϕ on completions

$$\hat{\phi} : \widehat{\mathscr{O}}_{\mathbb{A}^1_k \times_k \mathbb{A}^1_k,(0,0)} \to \widehat{\mathscr{O}}_{\mathbb{G}_{m,k} \times_k \mathbb{A}^1_k,(1,0)}$$

is an isomorphism. This homomorphism is given by

$$\begin{aligned} &k[[x, z']] \to k[[x - 1, z']], \\ &x \mapsto (x - 1)\delta(x - 1, z'), \quad z' \mapsto z'. \end{aligned}$$

Here we regard $\delta(x-1,z')$ as a square root of $\frac{g(x,z')}{(x-1)^2}$ in $k[[x-1,z']]$. Using the fact that the image of $\delta(x-1,z')$ in the residue field, that is, the constant term of $\delta(x-1,z')$, is nonzero, one can check the above homomorphism on formal power series rings is indeed an isomorphism. □

Lemma 1.4. *Let $g(x,z') \in k[x,1/x,z']$ such that*

$$\begin{aligned} &g(1,z') = 0, \\ &\frac{\partial g}{\partial x}(1,z') = 0, \\ &\frac{1}{2}\frac{\partial^2 g}{\partial x^2}(1,0) \text{ is a nonzero square in } k, \end{aligned}$$

Write $g(x,z')$ as

$$g(x,z') = \sum_{i=-m}^{n} c_i(z')x^i,$$

where $m,n \geq 0$ and $c_i(z') \in k[z']$. Let $\mathscr{K}$ be a rank 1 lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ such that $\mathscr{K}|_1 \cong \overline{\mathbb{Q}}_\ell$.

(i) Suppose $m,n \geq 1$ and $c_n(0)c_{-m}(0) \neq 0$. Denote by G the k-morphism

$$G : \mathbb{G}_{m,k} \times_k \mathbb{G}_{m,k} \to \mathbb{A}^1_k \times_k \mathbb{G}_{m,k}, \quad (x,t') \to (g(x,1/t'),t').$$

For any $\lambda \in k$, $\big(\mathscr{K}_{\chi_2}(\lambda t'^s) \otimes G(\chi_2,\psi)\big)|_{\eta'_\infty}$ is a direct factor of

$$\Big(R^1\mathrm{pr}_{2!}\big(RG_!(\pi_1^*\mathscr{K}_Q) \otimes \mathscr{L}_\psi(\lambda x t'^s)\big)\Big)|_{\eta_{\infty'}},$$

where

$$\pi_1 : \mathbb{G}_{m,k} \times_k \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}, \quad \mathrm{pr}_2 : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$$

are the projections to the first and second factors, respectively.

(ii) Suppose $m = 0$, $n \geq 1$ and $c_n(0) \neq 0$. Denote by G the k-morphism

$$G : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{A}^1_k \times_k G_{m,k}, \quad (x,t') \to (g(x,1/t'),t').$$

For any $\lambda \in k$, $\big(\mathscr{K}_{\chi_2}(\lambda t'^s) \otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$ is a direct factor of

$$\Big(R^1\mathrm{pr}_{2!}\big(RG_!(\mathrm{pr}_1^* j_! \mathscr{K}) \otimes \mathscr{L}_\psi(\lambda x t'^s)\big)\Big)|_{\eta_{\infty'}},$$

where $j : \mathbb{G}_{m,k} \hookrightarrow \mathbb{A}^1_k$ is the open immersion, and

$$\mathrm{pr}_1 : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{A}^1_k, \quad \mathrm{pr}_2 : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$$

are the projections.

Proof. We give a proof of (i). The morphism G can be extended to a k-morphism

$$\bar{G} : \mathbb{G}_{m,k} \times_k (\mathbb{G}_{m,k} \cup \{\infty'\}) \to \mathbb{A}^1_k \times_k (\mathbb{G}_{m,k} \cup \{\infty'\})$$

as follows: We have

$$\mathbb{G}_{m,k} \cup \{\infty'\} = \operatorname{Spec} k[z'].$$

The morphism $\bar{G}$ is

$$\bar{G} : \mathbb{G}_{m,k} \times_k (\mathbb{G}_{m,k} \cup \{\infty'\}) \to \mathbb{A}^1_k \times_k (\mathbb{G}_{m,k} \cup \{\infty'\}), \quad (x, z') \mapsto (g(x, z'), z').$$

Fix notations by the following commutative diagram:

$$\begin{array}{ccccccc}
\mathbb{G}_{m,k} & = & \mathbb{G}_{m,k} & = & \mathbb{G}_{m,k} & & \\
\uparrow \pi_1 & & \uparrow \bar{\pi}_1 & & \uparrow \tilde{\pi}_1 & & \\
\mathbb{G}_{m,k} \times_k \mathbb{G}_{m,k} & \hookrightarrow & \mathbb{G}_{m,k} \times_k (\mathbb{G}_{m,k} \cup \{\infty'\}) & \hookleftarrow & \mathbb{G}_{m,R} = \mathbb{G}_{m,k} \times_k \operatorname{Spec} R & \hookrightarrow & \mathbb{P}^1_R \\
\downarrow G & & \downarrow \bar{G} & & \downarrow \tilde{G} & & \downarrow \hat{G} \\
\mathbb{A}^1_k \times_k \mathbb{G}_{m,k} & \overset{\iota}{\hookrightarrow} & \mathbb{A}^1_k \times_k (\mathbb{G}_{m,k} \cup \{\infty'\}) & \hookleftarrow & \mathbb{A}^1_R = \mathbb{A}^1_k \times_k \operatorname{Spec} R & \overset{\kappa}{\hookrightarrow} & \mathbb{P}^1_R = \mathbb{P}^1_k \times_k \operatorname{Spec} R \\
\mathrm{pr}_2 \downarrow & & \downarrow \bar{\mathrm{pr}}_2 & & \downarrow \tilde{\mathrm{pr}}_2 & & \downarrow \hat{\mathrm{pr}}_2 \\
\mathbb{G}_{m,k} & \overset{\delta}{\hookrightarrow} & \mathbb{G}_{m,k} \cup \{\infty'\} = \operatorname{Spec} A & \hookleftarrow & \operatorname{Spec} R & = & \operatorname{Spec} R,
\end{array}$$

where $R = k[z']_{c_n c_{-m}}$ is the localization of $k[z']$ with respect to $c_n(z')c_{-m}(z')$, π_1, $\bar{\pi}_1$ and $\tilde{\pi}_1$ are projections to the first factors, pr_2, $\bar{\mathrm{pr}}_2$, $\tilde{\mathrm{pr}}_2$ and $\hat{\mathrm{pr}}_2$ are projections to the second factors, κ, ι and δ are the canonical open immersions. We have

$$\begin{aligned}
\Big(R\mathrm{pr}_{2!}\big(RG_!(\pi_1^*\mathscr{K}) \otimes \mathscr{L}_\psi(\lambda x t'^s)\big)\Big)|_{\eta_{\infty'}} &\cong \Big(R\bar{\mathrm{pr}}_{2!}\big(R\bar{G}_!(\bar{\pi}_1^*\mathscr{K}) \otimes \iota_!\mathscr{L}_\psi(\lambda x t'^s)\big)\Big)|_{\eta_{\infty'}} \\
&\cong \Big(R\tilde{\mathrm{pr}}_{2!}\big(R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K}) \otimes (\iota_!\mathscr{L}_\psi(\lambda x t'^s))|_{\mathbb{A}^1_R}\big)\Big)|_{\eta_{\infty'}} \\
&\cong \Big(R\hat{\mathrm{pr}}_{2*}\big(\kappa_! R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K}) \otimes \kappa_!((\iota_!\mathscr{L}_\psi(\lambda x t'^s))|_{\mathbb{A}^1_R})\big)\Big)|_{\eta_{\infty'}}.
\end{aligned}$$

Let $a_0(x) \in k[x, 1/x]$ be the constant term of $g(x, z') \in k[x, 1/x, z']$ consider as a polynomial of z'. Since $\frac{\partial^2 g}{\partial x^2}(1, 0)$ is nonzero, $\frac{\mathrm{d}a_0}{\mathrm{d}x}$ is nonzero. Let S be the set of those $a \in \bar{k}$ such that $a = a_0(b)$ for some $b \in \bar{k}$ satisfying $\frac{\mathrm{d}a_0}{\mathrm{d}x}(b) = 0$. Then S is a finite set. Since $g(1, z') = \frac{\partial g}{\partial x}(1, z') = 0$, we have $0 \in S$. By Lemma 1.2, $R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})$ is lisse in a neighborhood of (x, ∞') if $x \notin S$. By [18] 1.3.1.2, $\iota_!\mathscr{L}_\psi(xt')$ is universally strongly locally acyclic relative to $\bar{\mathrm{pr}}_2$. Since $\iota_!\mathscr{L}_\psi(\lambda x t'^s)$ is obtained from $\iota_!\mathscr{L}_\psi(xt')$ by base change, $\iota_!\mathscr{L}_\psi(\lambda x t'^s)$ is also universally strongly locally acyclic relative to $\bar{\mathrm{pr}}_2$. It follows that the vanishing cycle

$$R\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_! R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K}) \otimes \kappa_!((\iota_!\mathscr{L}_\psi(\lambda x t'^s))|_{\mathbb{A}^1_R})\big)$$

is supported at those points (x,∞') in $\mathbb{P}^1_k\times\infty'$ with $x=\infty$ or $x\in S$. Moreover, as $\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})$ vanishes on $\mathbb{P}^1_k\times\infty'$, its vanishing cycle and nearby cycle coincide, that is,

$$R\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\cong R\Psi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big).$$

So we have

$$\begin{aligned}
&\Big(R\hat{\mathrm{pr}}_{2*}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)_{\bar{\eta}_{\infty'}}\\
\cong\ & R\Gamma\Big(\mathbb{P}^1_{\bar{k}}\times\infty', R\Psi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)\\
\cong\ & R\Gamma\Big(\mathbb{P}^1_{\bar{k}}\times\infty', R\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)\\
\cong\ & \bigoplus_{x\in S}\Big(R\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)_{(x,\infty')}\\
&\bigoplus\Big(R\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)_{(\infty,\infty')}.
\end{aligned}$$

In particular, taking $x=0\in S$, we see

$$\Big(R\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)_{(0,\infty')}$$

is a direct factor of

$$\Big(R\hat{\mathrm{pr}}_{2*}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)_{\bar{\eta}_{\infty'}}.$$

We have

$$(\tilde{\pi}_1^*\mathscr{K})|_{(\mathbb{G}_{m,k}\times_k(\mathbb{G}_{m,k}\cup\{\infty'\}))^{\mathrm{h}}_{(1,\infty')}}\cong\overline{\mathbb{Q}}_l$$

since $\mathscr{K}$ is lisse on $\mathbb{G}_{m,k}$ and $\mathscr{K}|_1\cong\overline{\mathbb{Q}}_l$, where $(\mathbb{G}_{m,k}\times_k(\mathbb{G}_{m,k}\cup\{\infty'\}))^{\mathrm{h}}_{(1,\infty')}$ is the spectrum of the henselization of $\mathbb{G}_{m,k}\times_k(\mathbb{G}_{m,k}\cup\{\infty'\})$ at $(1,\infty')$. Let ω and ϕ be the morphisms defined in Lemma 1.3. Then $(R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K}))|_{(\mathbb{A}^1_k\times_k(\mathbb{G}_{m,k}\cup\{\infty'\}))^{\mathrm{h}}_{(0,\infty')}}$ has a direct factor $(\omega\circ\phi)_*(\overline{\mathbb{Q}}_l)$. Since ϕ is an isomorphism, we have $(\omega\circ\phi)_*(\overline{\mathbb{Q}}_l)\cong\omega_*(\overline{\mathbb{Q}}_l)$. But $\omega_*(\overline{\mathbb{Q}}_l)$ has a direct factor $(\tilde{\mathrm{pr}}_1^*j_!\mathscr{K}_{\chi_2})|_{(\mathbb{A}^1_k\times_k(\mathbb{G}_{m,k}\cup\{\infty'\}))^{\mathrm{h}}_{(0,\infty')}}$, where $\tilde{\mathrm{pr}}_1:\mathbb{A}^1_k\times_k\operatorname{Spec}R\to\mathbb{A}^1_k$ is the projection. So

$$\Big(R^1\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_!R\tilde{G}_!(\tilde{\pi}_1^*\mathscr{K})\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)_{(0,\infty')}$$

has a direct factor

$$\Big(R^1\Phi_{\bar{\eta}_{\infty'}}\big(\kappa_!\tilde{\mathrm{pr}}_1^*j_!\mathscr{K}_{\chi_2}\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda xt'^s))|_{\mathbb{A}^1_R})\big)\Big)_{(0,\infty')}.$$

Let $f:\eta_{\infty'}\to\eta_{\eta'}$ be the k-morphism defined by the k-algebra homomorphism

$$k((1/t'))\to((1/t')),\quad t'\mapsto\lambda t'^s.$$

Then we have

$$\begin{aligned}\Big(R^1\Phi_{\bar\eta_{\infty'}}\big(\kappa_!\tilde{\mathrm{pr}}_1^*j_!\mathscr{K}_{\chi_2}\otimes\kappa_!((\iota_!\mathscr{L}_\psi(\lambda x t'^s))|_{\mathbb{A}^1_R})\big)\Big)_{(0,\infty')} &\cong f^*\mathfrak{F}^{(0,\infty')}(\mathscr{K}_{\chi_2}|_{\eta_0})\\ &\cong f^*\big(\mathscr{K}_{\chi_2}\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}\\ &\cong \big(\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}},\end{aligned}$$

where for the first isomorphism, we use [5] Th. finitude 3.7 about the base change of vanishing cycles, and for the second isomorphism, we use [18] 2.5.3.1 (ii). This proves Lemma 1.4. □

In the rest of this section, we denote by $\pi_1,\pi_2:\mathbb{G}_{m,k}\times_k\mathbb{G}_{m,k}\to\mathbb{G}_{m,k}$ the two projections.

Lemma 1.5. *Let $\mathscr{K}$ be a rank 1 lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ such that $\mathscr{K}|_1\cong\overline{\mathbb{Q}}_\ell$, let $f(x,t')\in k[x,1/x,t',1/t']$, and let $a_s(x)t'^s$ be the highest degree term of $f(x,t')$ considered as a Laurent polynomial of t', where $a_s(x)\in k[x,1/x]$. Suppose*

$$\frac{\partial f}{\partial x}(1,t')=0,$$
$$\lambda=\frac{1}{2}\frac{\partial^2 a_s}{\partial x^2}\Big|_{x=1}\ \text{is nonzero.}$$

Write $f(x,t')$ as

$$f(x,t')=\sum_{i=-m}^{n}c_i(t')x^i,$$

where $m,n\geq 0$ and $c_i(t')\in k[t',1/t']$. Suppose one of the following conditions hold:

(a) $m,n\geq 1$ and $\Big(\frac{c_n(t')}{t'^s}\cdot\frac{c_{-m}(t')}{t'^s}\Big)|_{t'=\infty}\neq 0$.

(b) $m=0$, $n\geq 1$ and $\frac{c_n(t')}{t'^s}|_{t'=\infty}\neq 0$.

(c) $m\geq 1$, $n=0$ and $\frac{c_{-m}(t')}{t'^s}|_{t'=\infty}\neq 0$.

Then

$$\big(\mathscr{L}_\psi(f(1,t'))\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$$

is a direct factor of

$$\Big(R^1\pi_{2!}\big(\mathscr{L}_\psi(f(x,t'))\otimes{\pi_1}^*\mathscr{K}\big)\Big)|_{\eta_{\infty'}}.$$

Proof. Let

$$g(x,1/t')=\frac{f(x,t')-f(1,t')}{\lambda t'^s}$$

and set $z'=1/t'$. In the case (a), $g(x,z')$ satisfies the conditions of Lemma 1.4 (i). Let G be the k-morphism

$$G:\mathbb{G}_{m,k}\times_k\mathbb{G}_{m,k}\to\mathbb{A}^1_k\times_k\mathbb{G}_{m,k},\quad G(x,t')=(g(x,1/t'),t')$$

and let $\mathrm{pr}_2 : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$ be the projection. We have $\mathrm{pr}_2 \circ G = \pi_2$. Combined with the projection formula ([3] XVII 5.2.9), we get

$$\begin{aligned}
& R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t')) \otimes {\pi_1}^*\mathscr{K}\big) \\
\cong\ & R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t') - f(1,t')) \otimes \pi_2^*\mathscr{L}_\psi(f(1,t')) \otimes \pi_1^*\mathscr{K}\big) \\
\cong\ & \mathscr{L}_\psi(f(1,t')) \otimes R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t') - f(1,t')) \otimes \pi_1^*\mathscr{K}\big) \\
\cong\ & \mathscr{L}_\psi(f(1,t')) \otimes R(\mathrm{pr}_2 \circ G)_!\big(G^*\mathscr{L}_\psi(\lambda x t'^s) \otimes \pi_1^*\mathscr{K}\big) \\
\cong\ & \mathscr{L}_\psi(f(1,t')) \otimes R\mathrm{pr}_{2!}\big(\mathscr{L}_\psi(\lambda x t'^s) \otimes RG_!\pi_1^*\mathscr{K}\big).
\end{aligned}$$

Our assertion then follows from Lemma 1.4 (i).

In the case (b), $f(x,t')$ is a polynomial of x, and $g(x,z')$ satisfies the conditions of Lemma 1.4 (ii). Let G be the k-morphism

$$G : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{A}^1_k \times_k \mathbb{G}_{m,k}, \quad G(x,t') = (g(x,1/t'),t'),$$

let

$$\mathrm{pr}_1 : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{A}^1_k, \qquad \mathrm{pr}_2 : \mathbb{A}^1_k \times_k \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$$

be the projections, and let $j : \mathbb{G}_{m,k} \to \mathbb{A}^1_k$ be the open immersion. The same argument as above shows that

$$R\mathrm{pr}_{2!}\big(\mathscr{L}_\psi(f(x,t')) \otimes {\mathrm{pr}_1}^* j_!\mathscr{K}\big) \cong \mathscr{L}_\psi(f(1,t')) \otimes R\mathrm{pr}_{2!}\big(\mathscr{L}_\psi(\lambda x t'^s) \otimes RG_!\mathrm{pr}_1^* j_!\mathscr{K}\big).$$

Moreover, we have

$$\begin{aligned}
& R\mathrm{pr}_{2!}\big(\mathscr{L}_\psi(f(x,t')) \otimes {\mathrm{pr}_1}^* j_!\mathscr{K}\big) \\
\cong\ & R\mathrm{pr}_{2!}\big(\mathscr{L}_\psi(f(x,t')) \otimes (j \times \mathrm{id}_{\mathbb{G}_{m,k}})_!\pi_1^*\mathscr{K}\big) \\
\cong\ & R\mathrm{pr}_{2!}(j \times \mathrm{id}_{\mathbb{G}_{m,k}})_!\big((j \times \mathrm{id}_{\mathbb{G}_{m,k}})^*\mathscr{L}_\psi(f(x,t')) \otimes \pi_1^*\mathscr{K}\big) \\
\cong\ & R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t')) \otimes {\pi_1}^*\mathscr{K}\big).
\end{aligned}$$

So we have

$$R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t')) \otimes {\pi_1}^*\mathscr{K}\big) \cong \mathscr{L}_\psi(f(1,t')) \otimes R\mathrm{pr}_{2!}\big(\mathscr{L}_\psi(\lambda x t'^s) \otimes RG_!\mathrm{pr}_1^* j_!\mathscr{K}\big).$$

Our assertion then follows from Lemma 1.4 (ii).

In the case (c), $f(1/x,t')$ satisfies the condition (b). Our assertion follows from the case (b) and the fact that

$$R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t'))\otimes {\pi_1}^*\mathscr{K}\big)\cong R\pi_{2!}\big(\mathscr{L}_\psi(f(1/x,t'))\otimes {\pi_1}^*(\mathrm{inv}^*\mathscr{K})\big),$$

where $\mathrm{inv}:\mathbb{G}_{m,k}\to\mathbb{G}_{m,k}$ is the morphism $x\mapsto 1/x$. □

Lemma 1.6. *Let $\mathscr{K}$ be a rank 1 lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and ∞, and let P be the morphism*

$$P:\mathbb{G}_{m,k}\times_k\mathbb{G}_{m,k}\to\mathbb{G}_{m,k},\quad (x,t')\mapsto xt'.$$

Then there exists a lisse $\overline{\mathbb{Q}}_\ell$-sheaf $\mathscr{K}_P$ on $\mathbb{G}_{m,k}$ tamely ramified at 0 and ∞ such that

$$P^*\mathscr{K}\cong\pi_1^*\mathscr{K}_P\otimes\pi_2^*\mathscr{K},\quad \mathscr{K}_P|_1\cong\overline{\mathbb{Q}}_\ell.$$

Proof. Let $\mathscr{K}^{-1}=\mathscr{H}om(\mathscr{K},\overline{\mathbb{Q}}_\ell)$. We claim that the canonical morphism

$$\pi_1^*\pi_{1*}(P^*\mathscr{K}\otimes\pi_2^*\mathscr{K}^{-1})\to P^*\mathscr{K}\otimes\pi_2^*\mathscr{K}^{-1}$$

is an isomorphism. To prove this, we may make base extension from k to its algebraic closure $\bar{k}$. There exists a character $\chi:\varprojlim_{(N,p)=1}\mu_N(\bar{k})\to\overline{\mathbb{Q}}_\ell^*$ such that $\mathscr{K}|_{\mathbb{G}_{m,\bar{k}}}\cong\mathscr{K}_\chi$. By [5] Sommes trig. (1.3.1), we have $P^*\mathscr{K}_\chi\otimes\pi_2^*\mathscr{K}_\chi^{-1}\cong\pi_1^*\mathscr{K}_\chi$. We have

$$\pi_1^*\pi_{1*}(\pi_1^*\mathscr{K}_\chi)\cong\pi_1^*(\pi_{1*}\pi_1^*\mathscr{K}_\chi)\cong\pi_1^*(\mathscr{K}_\chi\otimes\pi_{1*}\pi_1^*\overline{\mathbb{Q}}_\ell)\cong\pi_1^*(\mathscr{K}_\chi\otimes\overline{\mathbb{Q}}_\ell)\cong\pi_1^*\mathscr{K}_\chi.$$

Hence the canonical morphism $\pi_1^*\pi_{1*}(\pi_1^*\mathscr{K}_\chi)\to\pi_1^*\mathscr{K}_\chi$ is an isomorphism. It follows that the canonical morphism $\pi_1^*\pi_{1*}(P^*\mathscr{K}\otimes\pi_2^*\mathscr{K}^{-1})\to P^*\mathscr{K}\otimes\pi_2^*\mathscr{K}^{-1}$ is an isomorphism over $\bar{k}$, and hence over k. Let $\mathscr{K}_P=\pi_{1*}(P^*\mathscr{K}\otimes\pi_2^*\mathscr{K}^{-1})$. Then $P^*\mathscr{K}\cong\pi_1^*\mathscr{K}_P\otimes\pi_2^*\mathscr{K}$. Taking the restriction at $(1,1)$ on both sides of this isomorphism, we get $\mathscr{K}_P|_1\cong\overline{\mathbb{Q}}_\ell$. □

Lemma 1.7. *Let $\alpha(t),\gamma(t),\delta(t)\in k[1,1/t]$, let $\gamma,\delta:\mathbb{G}_{m,k}\to\mathbb{A}^1_k$ be the k-morphisms defined by $\gamma(t),\delta(t)$, respectively, and let $\mathscr{K}$ be a $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$. Denote by $\mathfrak{F}:D^b_c(\mathbb{A}^1_k,\overline{\mathbb{Q}}_\ell)\to D^b_c(\mathbb{A}^1_k,\overline{\mathbb{Q}}_\ell)$ the global Fourier transformation. We have*

$$\delta^*\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\cong R\pi_{2!}\big(\mathscr{L}_\psi(\alpha(t)+\gamma(t)\delta(t'))\otimes\pi_1^*\mathscr{K}\big)[1].$$

Proof. Fix notations by the following commutative diagram:



Applying the proper base change theorem and the projection formula, we get

$$\begin{aligned}
& \delta^* R p_{2!}\Big(p_1^*\gamma_!\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big) \otimes \mathscr{L}_\psi(tt')\Big) \\
\cong\ & R\mathrm{pr}_{2!}(\mathrm{id}_{\mathbb{A}^1_k} \times \delta)^*\Big(p_1^*\gamma_!\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big) \otimes \mathscr{L}_\psi(tt')\Big) \\
\cong\ & R\mathrm{pr}_{2!}\Big(\big(p_1 \circ (\mathrm{id}_{\mathbb{A}^1_k} \times \delta)\big)^*\gamma_!\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big) \otimes (\mathrm{id}_{\mathbb{A}^1_k} \times \delta)^*\mathscr{L}_\psi(tt')\Big) \\
\cong\ & R\mathrm{pr}_{2!}\Big((\gamma \times \mathrm{id}_{\mathbb{G}_{m,k}})_!\pi_1^*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big) \otimes (\mathrm{id}_{\mathbb{A}^1_k} \times \delta)^*\mathscr{L}_\psi(tt')\Big) \\
\cong\ & R\mathrm{pr}_{2!}(\gamma \times \mathrm{id}_{\mathbb{G}_{m,k}})_!\Big(\pi_1^*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}\big) \otimes (\gamma \times \mathrm{id}_{\mathbb{G}_{m,k}})^*(\mathrm{id}_{\mathbb{A}^1_k} \times \delta)^*\mathscr{L}_\psi(tt')\Big) \\
\cong\ & R\pi_{2!}\big(\mathscr{L}_\psi(\alpha(t) + \gamma(t)\delta(t')) \otimes \pi_1^*\mathscr{K}\big).
\end{aligned}$$

Our assertion follows. □

Proposition 1.8. *Let $\mathscr{K}$ be a rank 1 lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞, and let $\alpha(t), \gamma(t) \in k[t, 1/t]$. Suppose $\frac{\mathrm{d}\gamma}{\mathrm{d}t}$ divides $\frac{\mathrm{d}\alpha}{\mathrm{d}t}$ in $k[t, 1/t]$, and let*

$$\begin{aligned}
\delta(t) &= -\frac{\mathrm{d}\alpha}{\mathrm{d}\gamma} = -\frac{\frac{\mathrm{d}\alpha}{dt}}{\frac{\mathrm{d}\gamma}{\mathrm{d}t}}, \\
\beta(t') &= \alpha(t') + \gamma(t')\delta(t').
\end{aligned}$$

Let $a_s t^s$ be the highest degree term of $\alpha(t)$, and let r be the highest degree of terms in $\gamma(t)$. Suppose $p = \mathrm{char}\, k$ is relatively prime to $2, r, s$ and $s - r$. Write

$$\alpha(xt') + \gamma(xt')\delta(t') = \sum_{i=-m}^{n} c_i(t')x^i$$

with $m, n \geq 0$ and $c_i(t') \in k[t', 1/t']$. Suppose furthermore that one of the following conditions holds:

(a) $m, n \geq 1$ and $\left(\frac{c_n(t')}{t'^s} \cdot \frac{c_{-m}(t')}{t'^s}\right)|_{t'=\infty} \neq 0$.

(b) $m = 0$, $n \geq 1$ and $\frac{c_n(t')}{t'^s}|_{t'=\infty} \neq 0$.

(c) $m \geq 1$, $n = 0$ and $\frac{c_{-m}(t')}{t'^s}|_{t'=\infty} \neq 0$.

Then

$$\left(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}\left(\frac{1}{2}s(s-r)a_s t'^s\right)\otimes G(\chi_2,\psi)\right)|_{\eta_{\infty'}}$$

is a direct factor of

$$\left(\delta^*\mathscr{H}^0\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\right)|_{\eta_{\infty'}}.$$

Proof. Let $\lambda=\frac{1}{2}s(s-r)a_s$. By Lemma 1.7, it suffices to show

$$\left(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\right)|_{\eta_{\infty'}}$$

is a direct factor of

$$R^1\pi_{2!}\big(\mathscr{L}_\psi(\alpha(t)+\gamma(t)\delta(t'))\otimes\pi_1^*\mathscr{K}\big)|_{\eta_{\infty'}}.$$

Introduce a change of variables

$$x=\frac{t}{t'}$$

and let

$$f(x,t')=\alpha(t)+\gamma(t)\delta(t')=\alpha(xt')+\gamma(xt')\delta(t').$$

Then we have

$$\beta(t')=f(1,t').$$

Consider the isomorphism

$$\tau:\mathbb{G}_{m,k}\times_k\mathbb{G}_{m,k}\to\mathbb{G}_{m,k}\times_k\mathbb{G}_{m,k},\quad (x,t')\mapsto(xt',t')\,.$$

We have $\pi_2\tau=\pi_2$. So we have

$$\begin{aligned}
R\pi_{2!}\big(\mathscr{L}_\psi(\alpha(t)+\gamma(t)\delta(t'))\otimes{\pi_1}^*\mathscr{K}\big) &\cong R\pi_{2!}\tau^*\big(\mathscr{L}_\psi(\alpha(t)+\gamma(t)\delta(t'))\otimes{\pi_1}^*\mathscr{K}\big)\\
&\cong R\pi_{2!}\big(\tau^*\mathscr{L}_\psi(\alpha(t)+\gamma(t)\delta(t'))\otimes\tau^*{\pi_1}^*\mathscr{K}\big)\\
&\cong R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t'))\otimes({\pi_1}^*\mathscr{K}_P\otimes{\pi_2}^*\mathscr{K})\big)\\
&\cong R\pi_{2!}\big(\mathscr{L}_\psi(f(x,t'))\otimes{\pi_1}^*\mathscr{K}_P\big)\otimes\mathscr{K}\,.
\end{aligned}$$

Here for third isomorphism, we use Lemma 1.6, and for the fourth isomorphism, we use the projection formula. We will verify shortly that the conditions of Lemma 1.5 hold for $f(x,t')$. We then conclude from lemma 1.5 that

$$\big(\mathscr{L}_\psi(f(1,t'))\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$$

is a direct factor of

$$\Big(R^1\pi_{2!}\big(\mathscr{L}_\psi(f(x,t'))\otimes {\pi_1}^*\mathscr{K}_P\big)\Big)|_{\eta_{\infty'}},$$

and hence

$$\big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$$

is a direct factor of

$$\Big(R^1\pi_{2!}\big(\mathscr{L}_\psi(\alpha(t)+\gamma(t)\delta(t'))\otimes {\pi_1}^*\mathscr{K}\big)\Big)|_{\eta_{\infty'}}.$$

This proves our assertion.

Let's verify $f(x,t')$ satisfies the conditions of Lemma 1.5. We have

$$\begin{aligned}\frac{\partial f}{\partial x} &= \frac{\partial}{\partial x}\Big(\alpha(xt')+\gamma(xt')\delta(t')\Big)\\ &= t'\frac{\mathrm{d}\alpha}{\mathrm{d}t}(xt')+t'\frac{\mathrm{d}\gamma}{\mathrm{d}t}(xt')\delta(t').\end{aligned}$$

Since $\delta(t)=-\frac{\frac{\mathrm{d}\alpha}{\mathrm{d}t}}{\frac{\mathrm{d}\gamma}{\mathrm{d}t}}$, we have

$$\frac{\partial f}{\partial x}(1,t')=t'\frac{\mathrm{d}}{\mathrm{d}t'}(\alpha(t'))+t'\frac{\mathrm{d}}{\mathrm{d}t'}(\gamma(t'))\delta(t')=0.$$

Write

$$\begin{aligned}\alpha(t) &= a_st^s+a_{s-1}t^{s-1}+\cdots,\\ \gamma(t) &= b_rt^r+b_{r-1}t^{r-1}+\cdots.\end{aligned}$$

We have

$$\begin{aligned}f(x,t') &= \alpha(xt')+\gamma(xt')\delta(t')\\ &= \alpha(xt')-\gamma(xt')\frac{\frac{\mathrm{d}}{\mathrm{d}t'}(\alpha(t'))}{\frac{\mathrm{d}}{\mathrm{d}t'}(\gamma(t'))}\\ &= a_s(xt')^s+a_{s-1}(xt')^{s-1}+\cdots\\ &\quad -\Big(b_r(xt')^r+b_{r-1}(xt')^{r-1}+\cdots\Big)\left(\frac{sa_st'^{s-1}+(s-1)a_{s-1}t'^{s-2}+\cdots}{rb_rt'^{r-1}+(r-1)b_{r-1}t'^{r-2}+\cdots}\right).\end{aligned}$$

From this expression, we see that the highest degree term of $f(x,t')$ considered as a Laurent polynomial of t' is $a_s(x)t'^s$ with

$$a_s(x)=a_sx^s-\frac{s}{r}a_sx^r.$$

Note that we have

$$\frac{1}{2}\frac{\partial^2}{\partial x^2}(a_s(x))|_{x=1}=\frac{1}{2}s(s-r)a_s=\lambda.$$

Since p is relatively prime to $2,r,s$ and $s-r$, we have $\frac{1}{2}\frac{\partial^2}{\partial x^2}(a_s(x))|_{x=1}\neq 0$. □

Remark 1.9. This remark gives a philosophical reason why we introduce the change of variable $x = \frac{t}{t'}$. By the facts on classical analysis recalled in the introduction, we expect a dominant part of

$$\Big(R^1\pi_{2!}\big(\mathscr{L}_\psi(\alpha(t)+\gamma(t)\delta(t'))\otimes {\pi_1}^*\mathscr{K}\big)\Big)|_{\eta_{\infty'}}$$

comes from the contribution on the curve

$$\frac{\partial}{\partial t}\Big(\alpha(t)+\gamma(t)\delta(t')\Big)=0.$$

By assumption, we have $\frac{\mathrm{d}\alpha}{\mathrm{d}t}+\frac{\mathrm{d}\gamma}{\mathrm{d}t}\delta(t)=0$. So $t=t'$ is a branch of the above curve. By the change of variable $x=\frac{t}{t'}$, we see $x=1$ is a branch of the curve. So a dominant part of the above cohomology should come from the contribution at $x=1$. This is indeed the case as we have seen.

In the next section, we will deduce Theorems 0.1-0.4 from the following lemma by a group theoretical argument:

Lemma 1.10. *Let $\gamma(t)=t^r$, let $\alpha(t)\in k[t,1/t]$, let*

$$\delta(t)=-\frac{\mathrm{d}\alpha}{\mathrm{d}\gamma}=-\frac{1}{rt^{r-1}}\frac{\mathrm{d}\alpha}{dt},$$
$$\beta(t')=\alpha(t')+\gamma(t')\delta(t'),$$

and let $\mathscr{K}$ be a rank 1 lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_{m,k}$ tamely ramified at 0 and at ∞. We have $\gamma(0)=0$ and $\gamma(\infty)=\infty$.

(i) Suppose $\alpha(t)$ is of the form

$$\alpha(t)=\frac{a_{-s}}{t^s}+\frac{a_{-(s-1)}}{t^{s-1}}+\cdots+\frac{a_{-1}}{t}$$

with $a_{-s}\neq 0$, and suppose $r,s\geq 1$ and $p=\operatorname{char}k$ is relatively prime to 2, r, s and $r+s$. We have $\delta(0')=\infty'$ and

$$\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}\left(\frac{1}{2}s(r+s)a_{-s}t'^s\right)\otimes G(\chi_2,\psi)\Big)|_{\eta_{0'}}$$

is a direct factor of $\delta^*\mathfrak{F}^{(0,\infty')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_0}\Big)$.

(ii) Suppose $\alpha(t)$ is of the form

$$\alpha(t)=a_st^s+a_{s-1}t^{s-1}+\cdots+a_1t$$

with $a_s\neq 0$, and suppose $s>r\geq 1$ and p is relatively prime 2, r, s and $s-r$. We have $\delta(\infty')=\infty'$ and

$$\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}\left(\frac{1}{2}s(s-r)a_st'^s\right)\otimes G(\chi_2,\psi)\Big)|_{\eta_{\infty'}}$$

is a direct factor of $\delta^*\mathfrak{F}^{(\infty,\infty')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_\infty}\Big)$.

(iii) Suppose $\alpha(t)$ *is of the form*

$$\alpha(t)=a_st^s+a_{s-1}t^{s-1}+\cdots+a_1t$$

with $a_s\neq 0$*, and suppose* $r>s\geq 1$ *and* p *is relatively prime to* 2*,* r*,* s *and* $s-r$*. We have* $\delta(\infty')=0'$ *and*

$$\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}\Big(\frac{1}{2}s(s-r)a_st'^s\Big)\otimes G(\chi_2,\psi)\Big)|_{\eta_{\infty'}}$$

is a direct factor of $\delta^*\mathfrak{F}^{(\infty,0')}\Big(\Big(\gamma_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_\infty}\Big)$.

Proof.

(i) Let

$$\begin{aligned}
&\gamma_1(t)=\gamma(1/t)=t^{-r},\\
&\alpha_1(t)=\alpha(1/t)=a_{-s}t^s+a_{-(s-1)}t^{s-1}+\cdots+a_{-1}t,\\
&\delta_1(t)=-\frac{\frac{\mathrm{d}}{\mathrm{d}t}(\alpha_1(t))}{\frac{\mathrm{d}}{\mathrm{d}t}(\gamma_1(t))}=\delta(1/t)=\frac{s}{r}a_{-s}t^{r+s}+\frac{s-1}{r}a_{-(s-1)}t^{r+s-1}+\cdots+\frac{1}{r}a_{-1}t^{r+1},\\
&\beta_1(t')=\alpha_1(t')+\gamma_1(t')\delta_1(t')=\beta(1/t'),
\end{aligned}$$

and let $\lambda=\frac{1}{2}s(r+s)a_{-s}$. From the above expression for $\delta_1(t')=\delta(1/t')$, we get $\delta(0')=\infty'$. Note that as a Laurent polynomial of x, the highest and the lowest degree terms of $\alpha_1(xt')+\gamma_1(xt')\delta_1(t')$ are $(a_{-s}t'^s)x^s$ and $\Big(\frac{s}{r}a_{-s}t'^s+\frac{s-1}{r}a_{-(s-1)}t'^{s-1}+\cdots+\frac{1}{r}a_{-1}t'\Big)x^{-r}$, respectively, and we have

$$\frac{a_{-s}t'^s}{t'^s}|_{t'=\infty'}\neq 0,\quad \frac{\frac{s}{r}a_{-s}t'^s+\frac{s-1}{r}a_{-(s-1)}t'^{s-1}+\cdots+\frac{1}{r}a_{-1}t'}{t'^s}|_{t'=\infty}\neq 0.$$

So we can apply Proposition 1.8 (a) to $\alpha_1(t)$, $\gamma_1(t)$ and $\mathrm{inv}^*\mathscr{K}$, and we see

$$\Big(\mathscr{L}_\psi(\beta_1(t'))\otimes\mathrm{inv}^*\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\Big)|_{\eta_{\infty'}}$$

is a direct factor of

$$\Big(\delta_1^*\mathscr{H}^0\mathfrak{F}\Big(\gamma_{1!}\big(\mathscr{L}_\psi(\alpha_1(t))\otimes\mathrm{inv}^*\mathscr{K}\big)\Big)\Big)|_{\eta_{\infty'}}.$$

We have

$$\begin{aligned}
\mathscr{L}_\psi(\beta_1(t'))\otimes\mathrm{inv}^*\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi) &\cong \mathrm{inv}^*\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\Big)\\
\gamma_{1!}\big(\mathscr{L}_\psi(\alpha_1(t))\otimes\mathrm{inv}^*\mathscr{K}\big) &\cong \gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big).
\end{aligned}$$

So

$$\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\Big)|_{\eta_{0'}}$$

is a direct factor of

$$\Big(\delta^*\mathscr{H}^0\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\Big)|_{\eta_{0'}}.$$

On the other hand, $\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)$ is tamely ramified at ∞ since $\alpha(t)$ is a polynomial of $1/t$ and $(r,p)=1$. By [18] 2.4.3 (iii) b) and the stationary phase principle [18] 2.3.3.1 (iii), as sheaves on $\eta_{\infty'}$, we have

$$\Big(\mathscr{H}^0\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\Big)|_{\eta_{\infty'}}\cong\mathfrak{F}^{(0,\infty')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_0}\Big).$$

So

$$\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\Big)|_{\eta_{0'}}$$

is a direct factor of $\delta^*\mathfrak{F}^{(0,\infty')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_0}\Big)$. Finally note that γ induces a finite morphism $\gamma:\mathbb{G}_{m,k}\to\mathbb{G}_{m,k}$. So we can replace $\gamma_!$ by γ_* in the above expression. Our assertion follows.

(ii) We have

$$\delta(t)=-\frac{1}{rt^{r-1}}\frac{\mathrm{d}\alpha}{\mathrm{d}t}=-\frac{s}{r}a_st^{s-r}-\frac{s-1}{r}a_{s-1}t^{s-r-1}-\cdots-\frac{1}{r}a_1t^{-r+1}.$$

Since $s>r$, we have $\delta(\infty')=\infty'$. Note that $\alpha(xt')+\gamma(xt')\delta(t')$ is a polynomial of x, and the highest degree term as a polynomial of x is $(a_st'^s)x^s$, and we have $\frac{a_st'^s}{t'^s}|_{t'=\infty'}\neq 0$. So we can apply Proposition 1.8 (b). Let $\lambda=\frac{1}{2}s(s-r)a_s$. Then

$$\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\Big)|_{\eta_{\infty'}}$$

is a direct factor of $\Big(\delta^*\mathscr{H}^0\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\Big)|_{\eta_{\infty'}}$. By the stationary phase principle [18] 2.3.3.1 (iii), as sheaves on $\eta_{\infty'}$, we have

$$\begin{aligned}&\Big(\mathscr{H}^0\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\Big)|_{\eta_{\infty'}}\\ \cong\ &\mathscr{F}^{(0,\infty')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_0}\Big)\bigoplus\mathscr{F}^{(\infty,\infty')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_\infty}\Big).\end{aligned}$$

Since $\alpha(t)$ is a polynomial of t, $\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_0}$ is tame and hence $\mathscr{F}^{(0,\infty')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_0}\Big)$ is also tame by [18] 2.5.3.1. But the slope of $\big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$ is

nonzero. So $\big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$ is a direct factor of $\delta^*\mathfrak{F}^{(\infty,\infty')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_\infty}\Big)$.

(iii) Again we have

$$\delta(t)=-\frac{1}{rt^{r-1}}\frac{\mathrm{d}}{\mathrm{d}t}(\alpha(t))=-\frac{s}{r}a_st^{s-r}-\frac{s-1}{r}a_{s-1}t^{s-r-1}-\cdots-\frac{1}{r}a_1t^{-r+1}.$$

Since $s<r$, we have $\delta(\infty')=0'$. Note that $\alpha(xt')+\gamma(xt')\delta_1(t')$ is a polynomial of x, and the highest degree term as a polynomial of x is

$$\Big(-\frac{s}{r}a_st'^s-\frac{s-1}{r}a_{s-1}t'^{s-1}-\cdots-\frac{1}{r}a_1t'\Big)x^r,$$

and we have

$$\frac{-\frac{s}{r}a_st'^s-\frac{s-1}{r}a_{s-1}t'^{s-1}-\cdots-\frac{1}{r}a_1t'}{t'^s}|_{t'=\infty}\neq 0.$$

So we can apply Proposition 1.8 (b). Let $\lambda=\frac{1}{2}s(s-r)a_s$. Then

$$\big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$$

is a direct factor of $\Big(\delta^*\mathscr{H}^0\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\Big)|_{\eta_{\infty'}}$. We have

$$\begin{aligned}\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)[1]\Big)_{\bar{0}} &\cong R\Gamma_c\big(\mathbb{A}^1_{\bar{k}},\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\big)[2]\\ &\cong R\Gamma_c\big(\mathbb{G}_{m,\bar{k}},\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)[2].\end{aligned}$$

Since the sheaf $\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}$ on $\mathbb{G}_{m,k}$ is nonconstant and lisse of rank 1, and thus irreducible, we have

$$H^2_c\big(\mathbb{G}_{m,\bar{k}},\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)=0$$

by [7] 1.4.1.3. So by [18] 2.3.2, we have a short exact sequence

$$\begin{aligned}0 &\to H^1_c\big(\mathbb{G}_{m,\bar{k}},\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\\ &\to \Big(\mathscr{H}^0\mathfrak{F}\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)\Big)|_{\bar{\eta}_{0'}}\\ &\to \mathscr{F}^{(\infty,0')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t'))\otimes\mathscr{K}\big)\Big)|_{\eta_\infty}\Big)\to 0.\end{aligned}$$

Note that $H^1_c\big(\mathbb{G}_{m,\bar{k}},\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)$ defines an unramified representation of $\mathrm{Gal}(\bar{\eta}_{0'}/\eta_{0'})$, whereas the representation defined by $\big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$ has no unramified subquotient. So $\big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}(\lambda t'^s)\otimes G(\chi_2,\psi)\big)|_{\eta_{\infty'}}$ is a direct factor of $\delta^*\mathfrak{F}^{(\infty,0')}\Big(\Big(\gamma_!\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)\Big)|_{\eta_\infty}\Big)$. □

2 Proof of Theorems

In this section, we use a group theoretical argument to deduce Theorems 0.1-0.4 from Lemma 1.10. We assume k is algebraically closed throughout this section except when we prove Theorem 0.1 in the end.

Lemma 2.1. *Let G be a group, let H be a subgroup of G, and let V be a finite dimensional representation of G over a field of characteristic* 0.

(i) If the restriction of this representation to H is semisimple and if H has finite index in G, then V is a semisimple representation of G.

(ii) If V is a semisimple representation of G and if H is a normal subgroup of G, then the restriction of this representation to H is semisimple.

Proof. (i) Let U be a subspace of V invariant under the action of G. We need to show that there exists a homomorphism $\pi : V \to U$ such that $\pi|_U$ is the identity and $g^{-1}\pi g = \pi$ for all $g \in G$. Since V is semisimple as a representation of H, we can find a homomorphism $\pi_0 : V \to U$ such that $\pi_0|_U$ is the identity and $h^{-1}\pi_0 h = \pi_0$ for all $h \in H$. Let $Hg_1, \ldots, Hg_n$ be a family of representatives of right cosets of H, where $n = [G : H]$. Define $\pi = \frac{1}{n}\sum_{i=1}^{n} g_i^{-1}\pi_0 g_i$. One can check π has the required property.

(ii) We may assume V is an irreducible representation of G. If U is an irreducible sub-representation of H contained in V, then we have $V = \sum_{g\in G} gU$ since the righthand side is invariant under G. Since H is normal in G, each gU is invariant under the action of H. So gU is a representation of H, and it is irreducible. As a sum of irreducible representation, V is semisimple as a representation of H. □

Fix a separable closure $\overline{k((t))}$ of the field $k((t))$ of Laurent series. Let $p = \operatorname{char} k$, let $I = \mathrm{Gal}\left(\overline{k((t))}/k((t))\right)$ and let P be the wild inertia subgroup. Then P is a profinite p-group, and we have a canonical isomorphism

$$I/P \overset{\cong}{\to} \varprojlim_{(N,p)=1} \mu_N(k), \quad \sigma \mapsto \left(\frac{\sigma(\sqrt[N]{t})}{\sqrt[N]{t}}\right).$$

A $\overline{\mathbb{Q}}_\ell$-representation of a topological group is always assumed to be finite dimensional and continuous. Recall that a representation is called indecomposable if it is not a direct sum of two proper sub-representations. Any finite dimensional representation is a direct sum of indecomposable repre-

sentations. Any $\overline{\mathbb{Q}}_\ell$-representation of I defines a $\overline{\mathbb{Q}}_\ell$-sheaf on $\eta_0 = \operatorname{Spec} k((t))$. Fix an element

$$\zeta = (\zeta_N) \in \varprojlim_{(N,p)=1} \mu_N(k)$$

so that ζ_N is a primitive N-th root of unity for each N prime to p. It is a topological generator of $\varprojlim_{(N,p)=1} \mu_N(k)$. Let $U(n)$ be the n-dimensional representation $\varprojlim_{(N,p)=1} \mu_N(k) \to \mathrm{GL}(n, \overline{\mathbb{Q}}_\ell)$ so that ζ acts through the unipotent $(n \times n)$-matrix with a single Jordan block

$$\begin{pmatrix} 1 & 1 & & \\ & 1 & \ddots & \\ & & \ddots & 1 \\ & & & 1 \end{pmatrix}.$$

Composed with the canonical projection

$$I \to I/P \cong \varprojlim_{(N,p)=1} \mu_N(k),$$

we get a $\overline{\mathbb{Q}}_\ell$-representation of I, which we still denote by $U(n)$. It is indecomposable. We denote the $\overline{\mathbb{Q}}_\ell$-sheaf on $\eta_0 = \operatorname{Spec} k((t))$ corresponding to this representation also by $U(n)$.

Fix an element $g \in I$ so that its image in $I/P \cong \varprojlim_{(N,p)=1} \mu_N(k)$ is ζ. Let J the inverse image of $\zeta^{\mathbb{Z}}$ in I. Then J is dense in I, and it is the subgroup of I generated by P and g. Define a topology on J so that open subgroups of P form a base of neighborhoods of the identity element in J. Then P is an open subgroup of J, and the restriction to J of the homomorphism $I \to \varprojlim_{(N,p)=1} \mu_N(k)$ induces an isomorphism of discrete groups

$$J/P \overset{\cong}{\to} \{\zeta^m | m \in \mathbb{Z}\} \cong \mathbb{Z}.$$

Let $\chi : \mathbb{Z} \to \overline{\mathbb{Q}}_\ell^*$ be a character. Denote by $\mathscr{K}_\chi$ the representation

$$J \to \overline{\mathbb{Q}}_\ell^*, \quad \sigma g^m \mapsto \chi(m) \text{ for any } \sigma \in P \text{ and } m \in \mathbb{Z}.$$

For any integer n, the restriction to J of the representation $U(n)$ of I is also denoted by $U(n)$.

Lemma 2.2. *(i) Let V be an indecomposable $\overline{\mathbb{Q}}_\ell$-representation of J. Then there exists an irreducible representation W of J and a positive integer n such that*

$$V \cong W \otimes U(n).$$

This factorization is unique up to isomorphism, and W is isomorphic to any irreducible sub-representation of V.

(ii) Let V be an irreducible $\overline{\mathbb{Q}}_\ell$-representation of J. Then there exists a character $\chi : \mathbb{Z} \to \overline{\mathbb{Q}}_\ell^*$ such that the representation $V \otimes \mathscr{K}_{\chi^{-1}}$ factors through a finite discrete quotient group of J.

(iii) If W is an irreducible $\overline{\mathbb{Q}}_\ell$-representation of J, then for any positive integer n, $W \otimes U(n)$ is indecomposable.

Proof. (i) Since V is a $\overline{\mathbb{Q}}_\ell$-representation and P is a profinite p-group, the restriction of this representation to P factors through a finite quotient of P. In particular, as a representation of P, V is semisimple. Let $\mathscr{S}$ be the set of isomorphic classes of irreducible representations of P which are sub-representations of V. Consider the isotypical decomposition

$$V = \bigoplus_{W \in \mathscr{S}} V_W$$

of V as a representation of P, where for each $W \in \mathscr{S}$, V_W is a direct sum of several copies of W. Since P is normal in J, for any $h \in J$, hV_W is invariant under the action of P. So as a representation of P, we also have

$$V = \bigoplus_{W \in \mathscr{S}} hV_W,$$

and each hV_W is isotypical (i.e. a direct sum of several copies of an irreducible representation). By the uniqueness of the isotypical decomposition, h induces a permutation of the set $\{V_W | W \in \mathscr{S}\}$. So J acts on this set. Since V is indecomposable as a representation of J, J acts transitively. Fix a representation $W \in \mathscr{S}$, and let

$$J' = \{h \in J | hV_W = V_W\}$$

be the stabilizer of V_W. Then J' has finite index in J (since the set $\{V_W | W \in \mathscr{S}\}$ is finite) and $P \subset J'$. Let $N = [J : J']$. Then J' is generated by P and g^N. The space V_W is invariant under the action of J' and hence defines a representation of J'. We have

$$V \cong \mathrm{Ind}_{J'}^{J}(V_W).$$

Let $\rho : J' \to \mathrm{GL}(V_W)$ be the homomorphism defined by the representation V_W of J', and let $\rho^{g^N} : J' \to \mathrm{GL}(V_W)$ be the homomorphism

$$\rho^{g^N}(\sigma) = \rho(g^{-N} \sigma g^N)$$

for any $\sigma \in J'$. Then $g^N : V_W \to V_W$ defines an isomorphism from the representation ρ^{g^N} to the representation ρ. But as a representation of P, V_W is a direct sum of several copies of the irreducible

representation W. By the uniqueness (up to isomorphism) of such decomposition, if $\rho_W : P \to \mathrm{GL}(W)$ is the homomorphism defined by the representation W of P, there exists an isomorphism $F : W \to W$ from the representation $\rho_W^{g^N}$ to the representation ρ_W, where $\rho_W^{g^N}$ is defined by

$$\rho_W^{g^N}(\sigma) = \rho_W(g^{-N}\sigma g^N)$$

for any $\sigma \in P$. Consider the map

$$\rho' : J' \to \mathrm{GL}(W), \quad \sigma g^{mN} \mapsto \rho_W(\sigma)F^m \text{ for any } \sigma \in P \text{ and any integer } m.$$

One can verify ρ' defines a representation of J'. Since W is an irreducible representation of P, ρ' defines an irreducible representation of J'. We have an isomorphism

$$W \otimes \mathrm{Hom}_P(W, V_W) \stackrel{\cong}{\to} V_W, \quad w \otimes f \mapsto f(w)$$

of representations of P, where P acts trivially on $\mathrm{Hom}_P(W, V_W)$. Define the action of $h \in J'$ on $f \in \mathrm{Hom}_P(W, V_W)$ to be $\rho(h) \circ f \circ \rho'(h^{-1})$. (Note that $\rho(h) \circ f \circ \rho'(h^{-1})$ is P-equivariant since P is normal in J'.) Then the above isomorphism is an isomorphism of representations of J'. Since P acts trivially on $\mathrm{Hom}_P(W, V_W)$, the action of J' on $\mathrm{Hom}_P(W, V_W)$ factors through J'/P. Since $V = \mathrm{Ind}_{J'}^{J}(V_W)$ is indecomposable, V_W is also indecomposable as a representation of J'. Hence $\mathrm{Hom}_P(W, V_W)$ is indecomposable as a representation of J'/P. But J'/P is a cyclic group of infinite order generated by the image of g^N. Therefore the Jordan form of g^N on $\mathrm{Hom}_P(W, V_W)$ has a single Jordan block. Let λ be the eigenvalue of g^N on $\mathrm{Hom}_P(W, V_W)$, and let K be the one dimensional representation of J'/P so that the image of g^N acts by the scalar multiplication by λ. Then as representations of J', we have

$$W \otimes \mathrm{Hom}_P(W, V_W) \cong (W \otimes K) \otimes (K^{-1} \otimes \mathrm{Hom}_P(W, V_W)).$$

Now $W \otimes K$ is still an irreducible representation of J', and $K^{-1} \otimes \mathrm{Hom}_P(W, V_W)$ is a representation of J'/P, and the Jordan form of g^N on $K^{-1} \otimes \mathrm{Hom}_P(W, V_W)$ has a single Jordan block with eigenvalue 1. Let n be the dimension of $\mathrm{Hom}_P(W, V_W)$ and let U be the $n \times n$ matrix

$$U = \begin{pmatrix} 1 & 1 & & \\ & 1 & \ddots & \\ & & \ddots & 1 \\ & & & 1 \end{pmatrix}.$$

One can verify

$$U^N = \begin{pmatrix} 1 & N & * & * \\ & 1 & \ddots & * \\ & & \ddots & N \\ & & & 1 \end{pmatrix}.$$

In particular, all eigenvalues of U^N are 1 and $U^N - I$ has rank $n-1$. So U^N has a single Jordan block. Hence as representations of J', we have

$$K^{-1} \otimes \mathrm{Hom}_P(W, V_W) \cong \mathrm{Res}^J_{J'}(U(n)).$$

We thus have

$$\begin{aligned} V &\cong \mathrm{Ind}^J_{J'}(V_W) \\ &\cong \mathrm{Ind}^J_{J'}\Big((W \otimes K) \otimes \mathrm{Res}^J_{J'}(U(n))\Big) \\ &\cong \mathrm{Ind}^J_{J'}(W \otimes K) \otimes U(n). \end{aligned}$$

We have seen that W is an irreducible representation of J'. Note that J' is a normal subgroup of J since $P \subset J' \subset J$ and J/P is abelian. Using this fact, the fact that $W \otimes K$ is an irreducible representation of J', and Proposition 22 in [21] §7.3, one can check $\mathrm{Res}^J_{J'}\mathrm{Ind}^J_{J'}(W \otimes K)$ is semisimple. By Lemma 2.1 (i), $\mathrm{Ind}^J_{J'}(W \otimes K)$ is a semisimple representation of J. As $V \cong \mathrm{Ind}^J_{J'}(W \otimes K) \otimes U(n)$ and V is indecomposable, $\mathrm{Ind}^J_{J'}(W \otimes K)$ is an irreducible representation of J. This finishes the proof of (i).

(ii) Keep the notations in the proof of (i). We have shown $V \cong \mathrm{Ind}^J_{J'}(V_W)$. Suppose V is irreducible. Then V_W is irreducible. We have shown $V_W \cong W \otimes \mathrm{Hom}_P(W, V_W)$. As V_W is irreducible and the action of g^N on $\mathrm{Hom}_P(W, V_W)$ has a single Jordan block, $\mathrm{Hom}_P(W, V_W)$ must have dimension 1. So after twisting W by a one dimensional representation, we may assume $V \cong \mathrm{Ind}^J_{J'}(W)$, where W is a representation of J' which is irreducible as a representation of P. Let $\rho : J' \to \mathrm{GL}(W)$ be the homomorphism defined by the representation W. $\rho(P)$ is a finite group. For any $h \in J'$, let $\rho(h)$ act on this finite group by conjugation. There exists a positive integer r such that the conjugation by $\rho(h^r)$ induces the identity map on $\rho(P)$. For example, we can take r to be the order of the automorphism group of $\rho(P)$. Thus we have

$$\rho(g^{rN})^{-1}\rho(\sigma)\rho(g^{rN}) = \rho(\sigma)$$

for any $\sigma \in P$, that is, $\rho(g^{rN})$ is a P-equivariant isomorphism of W. But W is an irreducible representation of P. So $\rho(g^{rN})$ is a scalar multiplication by Schur's Lemma. Choose $c \in \overline{\mathbb{Q}}_\ell$ such that

$\rho(g^{rN})$ is the scalar multiplication by c^{rN}. Define a character $\chi : \mathbb{Z} \to \overline{\mathbb{Q}}_\ell^*$ by $\chi(1) = c$. We claim that the representation $V \otimes \mathscr{K}_{\chi^{-1}}$ factors through a finite discrete quotient group of J. We have

$$V \otimes \mathscr{K}_{\chi^{-1}} \cong \mathrm{Ind}_{J'}^{J}(W) \otimes \mathscr{K}_{\chi^{-1}} \cong \mathrm{Ind}_{J'}^{J}\big(W \otimes \mathrm{Res}_{J'}^{J}(\mathscr{K}_{\chi^{-1}})\big).$$

It suffices to show $W \otimes \mathrm{Res}_{J'}^{J}\mathscr{K}_{\chi^{-1}}$ factors through a finite discrete quotient group of J'. By our construction, g^{rN} acts trivially on $W \otimes \mathrm{Res}_{J'}^{J}\mathscr{K}_{\chi^{-1}}$. Moreover, there exists an open subgroup P' of P such that P' acts trivially on $W \otimes \mathrm{Res}_{J'}^{J}\mathscr{K}_{\chi^{-1}}$. Since the subgroup of J' generated by P' and g^{rN} has finite index, our assertion follows.

(iii) Let $V = W \otimes U(n)$. We can find a filtration

$$V = F_0 \supset F_1 \supset \cdots \supset F_n = 0$$

of V by J-invariant subspaces such that $F_i/F_{i+1} \cong W$ for any $0 \le i \le n-1$. By the Jordan-Hölder theorem, any irreducible subquotient of V is isomorphic to W. Write

$$V = V_1 \oplus \cdots \oplus V_m$$

so that each V_i is indecomposable. By (i), we must have

$$V_i \cong W \otimes U(n_i)$$

for some positive integer n_i. We thus have

$$V = W \otimes U(n) \cong W \otimes (U(n_1) \oplus \cdots \oplus U(n_m)).$$

Since $U(n)$ is trivial as a representation of P, we have a canonical isomorphism

$$\begin{aligned}&\mathrm{Hom}_P(W, W) \otimes U(n) \to \mathrm{Hom}_P(W, W \otimes U(n)),\\ &f \otimes u \mapsto (w \mapsto f(w) \otimes u) \text{ for any } f \in \mathrm{Hom}_P(W, W),\ u \in U(n),\ w \in W.\end{aligned}$$

This isomorphism is J-equivariant. Through this isomorphism, the action of g on $\mathrm{Hom}_P(W, W \otimes U(n))$ is identified with

$$g|_{\mathrm{Hom}_P(W,W)} \otimes g|_{U(n)}.$$

Similarly, the action of g on $\mathrm{Hom}_P\Big(W, W \otimes (U(n_1) \oplus \cdots \oplus U(n_m))\Big)$ is identified with

$$g|_{\mathrm{Hom}_P(W,W)} \otimes (g|_{U(n_1)} \oplus \cdots \oplus g|_{U(n_m)}).$$

By (ii), there exists a character $\chi : \mathbb{Z} \to \overline{\mathbb{Q}}_\ell^*$ such that the representation $W \otimes \mathscr{K}_{\chi^{-1}}$ factors through a finite quotient of J. As representations of J, we have

$$\mathrm{Hom}_P(W, W) \cong \mathrm{Hom}_P(W \otimes \mathscr{K}_{\chi^{-1}}, W \otimes \mathscr{K}_{\chi^{-1}}).$$

It follows that J acts on $\mathrm{Hom}_P(W, W)$ through a finite quotient. So the matrix of g on $\mathrm{Hom}_P(W, W)$ is diagonalizable. Let $r = \dim(\mathrm{Hom}_P(W, W))$. Then $g|_{\mathrm{Hom}_P(W,W)} \otimes g|_{U(n)}$ has r Jordan blocks and each of them is of size $n \times n$. Similarly, $g|_{\mathrm{Hom}_P(W,W)} \otimes (g|_{U(n_1)} \oplus \cdots \oplus g|_{U(n_m)})$ has rm Jordan blocks. Among them, r blocks are of size $n_i \times n_i$ for each i. So we must have $m = 1$ and V is indecomposable. □

Corollary 2.3. *(i) Let V be an indecomposable $\overline{\mathbb{Q}}_\ell$-representation of I. Then there exist an irreducible representation W of I and a positive integer n such that*

$$V \cong W \otimes U(n).$$

This factorization is unique up to isomorphism, and W is isomorphic to any irreducible sub-representation of V.

(ii) Conversely, if W is an irreducible $\overline{\mathbb{Q}}_\ell$-representation of I, then for any positive integer n, $W \otimes U(n)$ is indecomposable.

Proof. (i) Let W be an irreducible representation of I contained in V. Since J is dense in I, V is also an indecomposable representation of J and W is an irreducible representation of J contained in V. By Lemma 2.2 (i), there exists a positive integer n such that we have $V \cong W \otimes U(n)$ as representations of J. Again because J is dense in I, we have $V \cong W \otimes U(n)$ as representations of I.

(ii) W is an irreducible representation of J. By Lemma 2.2 (iii), $W \otimes U(n)$ is indecomposable as a representation of J. It follows that it is indecomposable as a representation of I. □

Remark 2.4. I don't know whether the analogue of Lemma 2.2 (ii) holds for irreducible representations of I, that is, whether the following statement is correct: Let W be an irreducible $\overline{\mathbb{Q}}_\ell$-representation of I. Then there exists a character $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$ such that the representation $W \otimes \mathcal{K}_{\chi^{-1}}$ factors through a finite discrete quotient of I, where $\mathcal{K}_{\chi^{-1}}$ is the Kummer representation of I associated to the character χ^{-1}. This is true if $\dim W = 1$. Indeed, let $\rho : I \to \overline{\mathbb{Q}}_\ell^*$ be the homomorphism defined by this representation. Then $\rho(g)$ is an ℓ-adic unit. So there exists a character $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$

such that $\chi(\xi) = \rho(g)$. Then $\rho\chi^{-1}$ is trivial on g and on an open subgroup of P. So $\rho\chi^{-1}$ factors through a finite quotient of I.

Lemma 2.5. *Any indecomposable representation of* $\varprojlim_{(N,p)=1} \mu_N(k)$ *is of the form* $\mathscr{K}_\chi \otimes U(n)$ *for some character* $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$ *and some integer* n.

Proof. Suppose $\rho : \varprojlim_{(N,p)=1} \mu_N(k) \to \mathrm{GL}(n, \overline{\mathbb{Q}}_\ell)$ is an indecomposable n-dimensional representation. Then $\rho(\zeta)$ is a matrix with a single Jordan block. Let λ be the eigenvalue of $\rho(\zeta)$ and let $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$ be the character defined by $\chi(\zeta) = \lambda$. Then ρ is isomorphic to $\mathscr{K}_\chi \otimes U(n)$. □

Lemma 2.6. *Let* $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_l^*$ *be a character, let* $\alpha(t) \in k((t))$ *be a formal Laurent series, and let* r *be a positive integer prime to* p. *Suppose for any* r*-th root of unity* $\mu \neq 1$, *the difference* $\alpha(\mu t) - \alpha(t)$ *is not of the form* $\gamma^p - \gamma$ $\big(\gamma \in k((t))\big)$. *Then* $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)$ *is an irreducible* $\overline{\mathbb{Q}}_\ell$*-sheaf on* η_0, *and for any positive integer* n, $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0} \otimes U(n)\big)$ *is an indecomposable* $\overline{\mathbb{Q}}_\ell$*-sheaf on* η_0.

Proof. Choose $T \in \overline{k((t))}$ so that $T^r = t$. Let $\mathscr{K}'$ be the inverse image of $\mathscr{K}_\chi$ under the morphism $\mathrm{Spec} k((T)) \to \mathbb{G}_{m,k}$ corresponding to the k-algebra homomorphism

$$k[t, 1/t] \to k((T)), \quad t \mapsto T,$$

and let $f : \mathrm{Spec}\, k((T)) \to \mathrm{Spec}\, k((t))$ be the morphism induced by the inclusion $k((t)) \hookrightarrow k((T))$. Then the sheaf $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)$ can be identified with $f_*\big(\mathscr{L}_\psi(\alpha(T)) \otimes \mathscr{K}'\big)$. Let's prove $f_*\big(\mathscr{L}_\psi(\alpha(T)) \otimes \mathscr{K}'\big)$ is irreducible. Let $I = \mathrm{Gal}\big(\overline{k((t))}/k((t))\big)$ and let $I' = \mathrm{Gal}\big(\overline{k((t))}/k((T))\big)$. Note that I' is a normal subgroup of I, and we have a canonical isomorphism

$$I/I' \xrightarrow{\cong} \mu_r(k), \quad gI' \mapsto \frac{g(T)}{T}.$$

Let $\rho : I' \to \mathrm{GL}(1, \overline{\mathbb{Q}}_\ell)$ be the representation corresponding to the rank 1 sheaf $\mathscr{L}_\psi(\alpha(T)) \otimes \mathscr{K}'$ on $\mathrm{Spec}\ k((T))$. Then the representation of I corresponding to the sheaf $f_*\big(\mathscr{L}_\psi(\alpha(T)) \otimes \mathscr{K}'\big)$ is $\mathrm{Ind}_{I'}^I(\rho)$. By the Mackey's criterion (Proposition 23 in [21] §7.4), to prove $\mathrm{Ind}_{I'}^I(\rho)$ is irreducible, it suffices to show that for any $g \in I - I'$, the representation ρ and ρ^g are disjoint, where $\rho^g : I' \to \mathrm{GL}(1, \overline{\mathbb{Q}}_\ell)$ is the representation defined by

$$\rho^g(\sigma) = \rho(g^{-1}\sigma g).$$

Let $\mu = \frac{g(T)}{T}$ and let

$$\tilde{g} : \operatorname{Spec} k((T)) \to \operatorname{Spec} k((T))$$

be the isomorphism induced by g. Then the sheaf on $\operatorname{Spec} k((T))$ corresponding to the representation ρ^g is

$$\tilde{g}^*(\mathscr{L}_\psi(\alpha(T)) \otimes \mathscr{K}') \cong \mathscr{L}_\psi(\alpha(\mu T)) \otimes \tilde{g}^*\mathscr{K}'.$$

By our assumption, $\alpha(\mu T) - \alpha(T)$ is not of the form $\gamma^p - \gamma$ $\big(\gamma \in k((T))\big)$. So $\mathscr{L}_\psi(\alpha(\mu T)) \otimes (\mathscr{L}_\psi(\alpha(T)))^{-1}$ is nontrivial. This representation is wildly ramified. As $\mathscr{K}' \otimes (\tilde{g}^*\mathscr{K}')^{-1}$ is tamely ramified, we have

$$\mathscr{L}_\psi(\alpha(\mu T)) \otimes (\mathscr{L}_\psi(\alpha(T)))^{-1} \ncong \mathscr{K}' \otimes (\tilde{g}^*\mathscr{K}')^{-1}.$$

Therefore,

$$\mathscr{L}_\psi(\alpha(\mu T)) \otimes \tilde{g}^*\mathscr{K}' \ncong \mathscr{L}_\psi(\alpha(T)) \otimes \mathscr{K}'.$$

So the two representations ρ and ρ^g of degree 1 are disjoint. This proves $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)$ is irreducible. We have

$$\begin{aligned} [r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0} \otimes U(n)\big) &\cong [r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0} \otimes [r]^*U(n)\big) \\ &\cong [r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big) \otimes U(n), \end{aligned}$$

and by Corollary 2.3 (ii), $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big) \otimes U(n)$ is indecomposable. So $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0} \otimes U(n)\big)$ is indecomposable. □

Lemma 2.7. *Let $\alpha(t) = \sum_{i=-s}^{\infty} a_i t^i \in k((t))$ be a formal Laurent series. For any positive integer r, let*

$$G(\alpha(t), r) = \{\mu \in \mu_r(k) | \alpha(\mu t) - \alpha(t) = \gamma^p - \gamma \text{ for some } \gamma \in k((t))\}.$$

Then $G(\alpha(t), r)$ is a group. Let $\alpha^{(1)}(t) = \sum_{i=-s}^{-1} a_i t^i$. Then we have $G(\alpha(t), r) = G(\alpha^{(1)}(t), r)$.

Proof. Suppose $\mu_1, \mu_2 \in G(\alpha(t), r)$. Choose $\gamma_1(t), \gamma_2(t) \in k((t))$ such that

$$\begin{aligned} \alpha(\mu_1 t) - \alpha(t) &= \gamma_1(t)^p - \gamma_1(t), \\ \alpha(\mu_2 t) - \alpha(t) &= \gamma_2(t)^p - \gamma_2(t). \end{aligned}$$

We have

$$\begin{aligned} \alpha(\mu_1\mu_2^{-1}t) - \alpha(t) &= (\alpha(\mu_1\mu_2^{-1}t) - \alpha(\mu_2^{-1}t)) + (\alpha(\mu_2^{-1}t) - \alpha(t)) \\ &= (\gamma_1(\mu_2^{-1}t)^p - \gamma_1(\mu_2^{-1}t)) - (\gamma_2(\mu_2^{-1}t)^p - \gamma_2(\mu_2^{-1}t)) \\ &= (\gamma_1(\mu_2^{-1}t) - \gamma_2(\mu_2^{-1}t))^p - (\gamma_1(\mu_2^{-1}t) - \gamma_2(\mu_2^{-1}t)). \end{aligned}$$

Therefore $\mu_1\mu_2^{-1} \in G(\alpha(t), r)$. So $G(\alpha(t), r)$ is a group.

Let $\alpha^{(2)}(t) = \sum_{i=0}^{\infty} a_i t^i$. Then for any $\mu \in \mu_r(k)$, $\alpha^{(2)}(\mu t) - \alpha^{(2)}(t)$ is a formal power series. By the Hensel Lemma, we can find a formal power series $\delta \in k[[t]]$ such that

$$\alpha^{(2)}(\mu t) - \alpha^{(2)}(t) = \delta^p - \delta.$$

Then $\alpha(\mu t) - \alpha(t) = \gamma^p - \gamma$ if and only if $\alpha^{(1)}(\mu t) - \alpha^{(1)}(t) = (\gamma - \delta)^p - (\gamma - \delta)$. It follows that $G(\alpha(t), r) = G(\alpha^{(1)}(t), r)$. □

Lemma 2.8. *Let $\mathscr{K}$ be a tamely ramified $\overline{\mathbb{Q}}_\ell$-sheaf on $\eta_0 = \operatorname{Spec} k((t))$ and let $\theta(t) \in k[[t]]$ be a formal power series of the form*

$$\theta(t) = a_1 t + a_2 t^2 + \cdots$$

with $a_1 \neq 0$. Denote by $\theta : \operatorname{Spec} k((t)) \to \operatorname{Spec} k((t))$ the morphism corresponding to the k-algebra homomorphism

$$k((t)) \to k((t)),\ t \mapsto \theta(t).$$

Then we have an isomorphism $\theta^*\mathscr{K} \cong \mathscr{K}$.

Proof. Set

$$\theta_1(t) = \frac{\theta(t)}{t} = a_1 + a_2 t + \cdots.$$

For each integer N with $(N, p) = 1$, choose an N-th root $\sqrt[N]{t}$ of t and an N-th root $\sqrt[N]{\theta_1(t)}$ of $\theta_1(t)$ in $\overline{k((t))}$. By the Hensel Lemma, we have $\sqrt[N]{\theta_1(t)} \in k[[t]]$ for all N. We can find a continuous representation $\rho : \varprojlim_{(N,p)=1} \mu_N(k) \to \mathrm{GL}(n, \overline{\mathbb{Q}}_\ell^*)$ so that $\mathscr{K}$ corresponds to the Galois representation

$$\mathrm{Gal}\left(\overline{k((t))}/k((t))\right) \to \mathrm{GL}(n, \overline{\mathbb{Q}}_\ell^*), \quad \sigma \mapsto \rho\left(\left(\frac{\sigma(\sqrt[N]{t})}{\sqrt[N]{t}}\right)_{(N,p)=1}\right).$$

Then $\theta^*\mathscr{K}$ corresponds to the Galois representation

$$\mathrm{Gal}\left(\overline{k((t))}/k((t))\right) \to \mathrm{GL}(n, \overline{\mathbb{Q}}_\ell^*), \quad \sigma \mapsto \rho\left(\left(\frac{\sigma(\sqrt[N]{t}\,\sqrt[N]{\theta_1(t)})}{\sqrt[N]{t}\,\sqrt[N]{\theta_1(t)}}\right)_{(N,p)=1}\right).$$

Since $\sqrt[N]{\theta_1(t)} \in k[[t]]$, we have $\sigma(\sqrt[N]{\theta_1(t)}) = \sqrt[N]{\theta_1(t)}$. Hence these two representations are the same.

□

Lemma 2.9. *Let $\chi : \varprojlim_{(N,p)=1} \mu_n(k) \to \overline{\mathbb{Q}}_\ell^*$ be a character. In the notation of Theorem 0.2, as sheaves on $\eta_{\infty'}$,*

$$\mathscr{L}_\psi(\beta(1/t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}$$

is a direct factor of $[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}_\chi|_{\eta_0}\big)\Big)$.

Proof. The equation

$$\frac{\mathrm{d}}{\mathrm{d}t}(\alpha(t))+rt^{r-1}t'^{r+s}=0$$

can be written as

$$t'^{r+s}=\frac{1}{t^{r+s}}\left(\frac{sa_{-s}}{r}+\frac{(s-1)a_{-(s-1)}}{r}t+\cdots\right).$$

Solve this equation, and write the solution as

$$t'=\frac{1}{t}(\lambda_0+\lambda_1 t+\cdots)=\lambda(t)$$

with $\lambda_0=\sqrt[r+s]{\frac{sa_{-s}}{r}}$. The solution is not unique and different solutions differ by an $(r+s)$-th root of unity. As long as λ_0 is fixed to be an $(r+s)$-th root of $\frac{sa_{-s}}{r}$, for each i, λ_i depends only on $a_{-s},a_{-(s-1)},\ldots,a_{-s+i}$. We can write

$$1/t'=\frac{t}{\lambda_0+\lambda_1 t+\cdots}=\nu_1 t+\nu_2 t^2+\cdots$$

with $\nu_1=\frac{1}{\lambda_0}$. For for each i, ν_i depends only on $\lambda_0,\ldots,\lambda_{i-1}$. Express t in terms of $1/t'$ and write the solution as

$$t=\mu_1(1/t')+\mu_2(1/t')^2+\cdots=\mu(1/t')$$

with $\mu_1=\frac{1}{\nu_1}=\sqrt[r+s]{\frac{sa_{-s}}{r}}$. For each i, μ_i depends only on $\nu_1,\ldots,\nu_i$. Substituting this expression into the equation

$$\beta(1/t')=\alpha(t)+t^r t'^{r+s},$$

we get

$$\begin{aligned}\beta(1/t') &= \alpha(\mu(1/t'))+(\mu(1/t'))^r t'^{r+s}\\ &= b_{-s}t'^s+b_{-(s-1)}t'^{s-1}+\cdots\end{aligned}$$

with

$$b_{-s}=\frac{a_{-s}}{\mu_1^s}+\mu_1^r=\frac{a_{-s}(1+\frac{s}{r})}{\left(\sqrt[r+s]{\frac{sa_{-s}}{r}}\right)^s}.$$

In particular, we have $b_{-s}\neq 0$. Note that in the above discussion, we need the fact that p is relatively prime to r, s and $r+s$. Moreover, as long as we fix an $(r+s)$-th root $\sqrt[r+s]{\frac{sa_{-s}}{r}}$ of $\frac{sa_{-s}}{r}$ and let

$b_{-s} = \frac{a_{-s}(1+\frac{s}{r})}{\left(\sqrt[r+s]{\frac{sa_{-s}}{r}}\right)^s}$, then for each i, b_i depends only on $a_{-s}, a_{-(s-1)}, \ldots, a_i$. If ζ is an $(r+s)$-th root of unity, then

$$t = \mu(1/\zeta t'),\ \beta = \beta(1/\zeta t')$$

is also a solution of the system of equations

$$\begin{cases} \alpha(t) + t^r t'^{r+s} = \beta(1/t'), \\ \frac{\mathrm{d}}{\mathrm{d}t}(\alpha(t)) + r t^{r-1} t'^{r+s} = 0, \end{cases}$$

and all the solutions of this system of equations are of this form. Note that

$$[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big)$$

does not depend on the choice of different solutions of β.

Write $\alpha(t) = \alpha^{(1)}(t) + \alpha^{(2)}(t)$, where

$$\alpha^{(1)}(t) = \sum_{i=-s}^{-1} a_i t^i, \quad \alpha^{(2)}(t) = \sum_{i=0}^{\infty} a_i t^i.$$

We have $\mathscr{L}_\psi(\alpha(t)) \cong \mathscr{L}_\psi(\alpha^{(1)}(t)) \otimes \mathscr{L}_\psi(\alpha^{(2)}(t))$, and$\mathscr{L}_\psi(\alpha^{(2)}(t)) \cong \overline{\mathbb{Q}}_\ell$ since k is algebraically closed. So we have

$$\mathscr{L}_\psi(\alpha(t)) \cong \mathscr{L}_\psi(\alpha^{(1)}(t)).$$

Write $\beta(1/t') = \beta^{(1)}(1/t') + \beta^{(2)}(1/t')$, where

$$\beta^{(1)}(1/t') = \sum_{i=-s}^{-1} b_i (1/t')^i, \quad \beta^{(2)}(1/t') = \sum_{i=0}^{\infty} b_i (1/t')^i.$$

We have

$$\mathscr{L}_\psi(\beta(1/t')) \cong \mathscr{L}_\psi(\beta^{(1)}(1/t')).$$

Note that $\beta^{(1)}(1/t')$ only depends on $\alpha^{(1)}(t)$. So to prove lemma, we may assume $\alpha(t)$ is of the form

$$\alpha(t) = \frac{a_{-s}}{t^s} + \frac{a_{-(s-1)}}{t^{s-1}} + \cdots + \frac{a_{-1}}{t}.$$

Let

$$\delta(t') = -\frac{1}{r t'^{r-1}} \frac{\mathrm{d}}{\mathrm{d}t'}(\alpha(t')) = \lambda(t')^{r+s},$$
$$\beta_0(t') = \alpha(t') + t'^r \lambda(t')^{r+s}.$$

Note that $\beta_0(t')$ is the $\beta(t')$ defined Theorem 0.1. We have

$$\begin{aligned}
\beta_0(\mu(1/t')) &= \alpha(\mu(1/t')) + (\mu(1/t'))^r \lambda(\mu(1/t'))^{r+s} \\
&= \alpha(\mu(1/t')) + (\mu(1/t'))^r t'^{r+s} \\
&= \beta(1/t'), \\
\delta(\mu(1/t')) &= \lambda(\mu(1/t'))^{r+s} \\
&= t'^{r+s}.
\end{aligned}$$

Let $\mu : \eta_{\infty'} \to \eta_{0'}$ and $\delta : \eta_{0'} \to \eta_{\infty'}$ be the k-morphisms corresponding to the k-algebra homomorphisms

$$\begin{aligned}
k((t')) \to k((1/t')), \qquad t' \mapsto \mu(1/t'), \\
k((1/t')) \to k((t')), \qquad t' \mapsto \delta(t'),
\end{aligned}$$

respectively. Then μ is an isomorphism and $\delta \circ \mu = [r+s]$. By Lemma 1.10 (i),

$$\left(\mathscr{L}_\psi(\beta_0(t')) \otimes \mathscr{K}_\chi \otimes \mathscr{K}_{\chi_2}\left(\frac{1}{2}s(r+s)a_{-s}t'^s\right) \otimes G(\chi_2, \psi)\right)|_{\eta_{0'}}$$

is a direct factor of $\delta^* \mathfrak{F}^{(0,\infty')}\left([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi\big)|_{\eta_0}\right)$. So

$$\mu^*\left(\left(\mathscr{L}_\psi(\beta_0(t')) \otimes \mathscr{K}_\chi \otimes \mathscr{K}_{\chi_2}\Big(\frac{1}{2}s(r+s)a_{-s}t'^s\Big) \otimes G(\chi_2, \psi)\Big)|_{\eta_{0'}}\right)$$

is a direct factor of $\mu^*\delta^* \mathfrak{F}^{(0,\infty')}\left([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi\big)|_{\eta_0}\right)$. By Lemma 2.8, we have

$$\begin{aligned}
\mu^* \mathscr{K}_\chi &\cong \mathscr{K}_{\chi^{-1}}, \\
\mu^* \mathscr{K}_{\chi_2}\left(\frac{1}{2}s(r+s)a_{-s}t'^s\right) &\cong [s]^* \mathscr{K}_{\chi_2}.
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
\mu^* \mathscr{L}_\psi(\beta_0(t')) &\cong \mathscr{L}_\psi(\beta_0(\mu(1/t'))) \\
&\cong \mathscr{L}_\psi(\beta(1/t')), \\
\mu^*\delta^* \mathfrak{F}^{(0,\infty')}\left([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi\big)|_{\eta_0}\right) &\cong (\delta \circ \mu)^* \mathfrak{F}^{(0,\infty')}\left([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi\big)|_{\eta_0}\right) \\
&\cong [r+s]^* \mathfrak{F}^{(0,\infty')}\left([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi\big)|_{\eta_0}\right)
\end{aligned}$$

Since k is algebraically closed, we have $G(\chi_2, \psi) \cong \overline{\mathbb{Q}}_\ell$. It follows that

$$\mathscr{L}_\psi(\beta(1/t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^* \mathscr{K}_{\chi_2})|_{\eta_{\infty'}}$$

is a direct factor of $[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}_\chi|_{\eta_0}\big)\Big)$. □

Lemma 2.10. *Let $\alpha(t)=\sum_{i=-s}^{\infty}a_it^i$ be a formal Laurent series in $k((t))$ and let $\alpha^{(1)}(t)=\sum_{i=-s}^{-1}a_it^i$. Define $\beta(1/t')=\sum_{i=-s}^{\infty}b_i(1/t')^i$ by the system of equations in Theorem 0.2.*

(i) If $\mu_d(k)\subset G(\alpha(t),r)$, then $d|r$, $d|s$ and $a_i=0$ for any i such that $-s\le i\le -1$ and $d\not|i$.

(ii) Under the assumption of (i), let $r_0=\frac{r}{d}$, $s_0=\frac{s}{d}$, $\alpha_0(t)=\sum_{d|i,\ i<0}a_it^{\frac{i}{d}}$, and let $\beta_0(1/t')$ be defined by the system of equations

$$\begin{cases}\alpha_0(t)+t^{r_0}t'^{r_0+s_0}=\beta_0(1/t'),\\ \frac{\mathrm{d}}{\mathrm{d}t}(\alpha_0(t))+r_0t^{r_0-1}t'^{r_0+s_0}=0.\end{cases}$$

Then $\beta(1/t')-\beta_0(1/t'^d)\in k[[t]]$.

(iii) $\mu_d(k)\subset G(\alpha(t),r)$ if and only if $\mu_d(k)\subset G(\beta(1/t'),r+s)$.

Proof. (i) Since $G(\alpha(t),r)$ is a subgroup of $\mu_r(k)$, we must have $d|r$. Fix a primitive d-th root of unity μ_0 and choose $\gamma\in k((t))$ so that

$$\alpha(\mu_0t)-\alpha(t)=\gamma^p-\gamma.$$

If the order $\mathrm{ord}_t(\gamma)$ of γ with respect to t is negative, that is, the formal Laurent series γ involves negative powers of t, then we have

$$\mathrm{ord}_t(\gamma^p-\gamma)=p\,\mathrm{ord}_t(\gamma)\le -p.$$

But

$$\mathrm{ord}_t(\alpha(\mu_0t)-\alpha(t))\ge -s.$$

So we have $-s\le -p$. This contradicts our assumption that $s<p$. So γ must be a formal power series, and hence the polar part

$$\sum_{i=-s}^{-1}a_i(\mu_0^i-1)t^i$$

of $\alpha(\mu_0t)-\alpha(t)$ vanishes. Thus $a_i=0$ whenever $i\le -1$ and $d\not|i$. As $a_{-s}\neq 0$, we have $d|s$.

(ii) Suppose we obtain from the equation

$$\frac{\mathrm{d}}{\mathrm{d}t}(\alpha_0(t))+r_0t^{r_0-1}t'^{r_0+s_0}=0$$

the expressions

$$t'=\lambda(t),\ t=\mu(1/t').$$

Then we have

$$\frac{\mathrm{d}}{\mathrm{d}t}(\alpha_0(t)) + r_0 t^{r_0-1}\lambda(t)^{r_0+s_0} = 0,$$
$$\beta_0(1/t') = \alpha_0(\mu(1/t')) + (\mu(1/t'))^{r_0} t'^{r_0+s_0}.$$

So we have

$$\begin{aligned} & \frac{\mathrm{d}}{\mathrm{d}t}(\alpha^{(1)}(t)) = \frac{\mathrm{d}}{\mathrm{d}t}(\alpha_0(t^d)) = \frac{\mathrm{d}\alpha_0}{\mathrm{d}t}(t^d)dt^{d-1} \\ = \quad & -r_0(t^d)^{r_0-1}\lambda(t^d)^{r_0+s_0}dt^{d-1} = -rt^{r-1}((\lambda(t^d))^{1/d})^{r+s}. \end{aligned}$$

Hence from the equation

$$\frac{\mathrm{d}}{\mathrm{d}t}(\alpha^{(1)}(t)) + rt^{r-1}t'^{r+s} = 0,$$

we get the expressions

$$t' = (\lambda(t^d))^{1/d},\ t = (\mu(1/t'^d))^{1/d},$$

and we have

$$\begin{aligned} \beta_0(1/t'^d) &= \alpha_0(\mu(1/t'^d)) + (\mu(1/t'^d))^{r_0}(t'^d)^{r_0+s_0} \\ &= \alpha^{(1)}((\mu(1/t'^d))^{1/d}) + ((\mu(1/t'^d))^{1/d})^r t'^{r+s}. \end{aligned}$$

So $\alpha^{(1)}(t)$ and $\beta_0(1/t'^d)$ satisfy the same system of equations for $\alpha(t)$ and $\beta(1/t')$:

$$\begin{cases} \alpha^{(1)}(t) + t^r t'^{r+s} = \beta_0(1/t'^d), \\ \frac{\mathrm{d}}{\mathrm{d}t}(\alpha^{(1)}(t)) + rt^{r-1}t'^{r+s} = 0. \end{cases}$$

Since $\alpha^{(1)}(t)$ is the polar part of $\alpha(t)$, $\beta(1/t)$ and $\beta_0(1/t'^d)$ have the same polar part by the discussion in the proof of Lemma 2.9. Hence $\beta(1/t) - \beta_0(1/t'^d) \in k[[t]]$.

(iii) Suppose $\mu_d(k) \subset G(\alpha(t), r)$. Then the conclusion of (ii) holds. It is clear that $\mu_d(k) \subset G(\beta_0(1/t'^d), r+s)$. Since $\beta(1/t') - \beta_0(1/t'^d) \in k[[t]]$, we have

$$G(\beta(1/t'), r+s) = G(\beta_0(1/t'^d), r+s)$$

by Lemma 2.7. So $\mu_d(k) \subset G(\beta(1/t'), r+s)$.

We have

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t'}(\beta(1/t')) &= \frac{\mathrm{d}}{\mathrm{d}t'}(\alpha(t) + t^r t'^{r+s}) \\ &= \frac{\mathrm{d}}{\mathrm{d}t}(\alpha(t))\frac{\mathrm{d}t}{\mathrm{d}t'} + rt^{r-1}t'^{r+s}\frac{\mathrm{d}t}{\mathrm{d}t'} + (r+s)t^r t'^{r+s-1} \\ &= \left(\frac{\mathrm{d}}{\mathrm{d}t}(\alpha(t)) + rt^{r-1}t'^{r+s}\right)\frac{\mathrm{d}t}{\mathrm{d}t'} + (r+s)t^r t'^{r+s-1} \\ &= (r+s)t^r t'^{r+s-1}. \end{aligned}$$

So $\alpha(t)$ is a solution of the system of equations

$$\begin{cases} \beta(1/t') - t^r t'^{r+s} = \alpha(t), \\ \frac{\mathrm{d}}{\mathrm{d}t'}(\beta(1/t')) - (r+s)t^r t'^{r+s-1} = 0. \end{cases}$$

(The above discussion is nothing but a re-proof of the involutivity of the Legendre transformation. See [2] Chapter 3, §14 C.) Reversing the role of α and β in the above discussion, we see that the condition $\mu_d(k) \subset G(\beta(1/t'), r+s)$ implies the condition $\mu_d(k) \subset G(\alpha(t), r)$. □

We are now ready to prove Theorems 0.1-0.4.

Proof of Theorem 0.2. Notation as in Theorem 0.2. We may assume $\alpha(t)$ is of the form

$$\alpha(t) = \frac{a_{-s}}{t^s} + \frac{a_{-(s-1)}}{t^{s-1}} + \cdots + \frac{a_{-1}}{t}$$

with $a_{-s} \neq 0$. First we prove that the general case of Theorem 0.2 follows from the special case where $\alpha(t)$ is assumed to have the property $G(\alpha(t), r) = 1$. Let $G(\alpha(t), r) = \mu_d(k)$ and let $r_0, s_0, \alpha_0(t), \beta_0(1/t')$ be defined as in Lemma 2.10. We have $\alpha(t) = \alpha_0(t^d)$, $G(\alpha_0(t), r_0) = \{1\}$, and $\beta(1/t') = \beta_0(1/t'^d)$. Note that $[d]_*\mathscr{K}$ is a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on $\mathbb{G}_m$ tamely ramified at 0 and at ∞. One can verify $\mathrm{inv}^*[d]_*\mathscr{K} \cong [d]_*\mathrm{inv}^*\mathscr{K}$. Applying the special case of Theorem 0.2 to $\alpha_0(t)$, we get

$$\begin{aligned} & \mathfrak{F}^{(0,\infty')}\Big([r_0]_*\big(\mathscr{L}_\psi(\alpha_0(t)) \otimes ([d]_*\mathscr{K})|_{\eta_0}\big)\Big) \\ \cong\ & [r_0+s_0]_*\big(\mathscr{L}_\psi(\beta_0(1/t')) \otimes (\mathrm{inv}^*[d]_*\mathscr{K})|_{\eta_{\infty'}} \otimes ([s_0]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big) \\ \cong\ & [r_0+s_0]_*\big(\mathscr{L}_\psi(\beta_0(1/t')) \otimes ([d]_*\mathrm{inv}^*\mathscr{K})|_{\eta_{\infty'}} \otimes ([s_0]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big). \end{aligned}$$

So we have

$$\begin{aligned} & \mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}|_{\eta_0}\big)\Big) \\ \cong\ & \mathscr{F}^{(0,\infty')}\Big([r_0]_*[d]_*\big([d]^*\mathscr{L}_\psi(\alpha_0(t)) \otimes \mathscr{K}|_{\eta_0}\big)\Big) \\ \cong\ & \mathfrak{F}^{(0,\infty')}\Big([r_0]_*\big(\mathscr{L}_\psi(\alpha_0(t)) \otimes ([d]_*\mathscr{K})|_{\eta_0}\big)\Big) \\ \cong\ & [r_0+s_0]_*\big(\mathscr{L}_\psi(\beta_0(1/t')) \otimes ([d]_*\mathrm{inv}^*\mathscr{K})|_{\eta_{\infty'}} \otimes ([s_0]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big) \\ \cong\ & [r_0+s_0]_*[d]_*\big([d]^*\mathscr{L}_\psi(\beta_0(1/t')) \otimes (\mathrm{inv}^*\mathscr{K})|_{\eta_{\infty'}} \otimes ([d]^*[s_0]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big) \\ \cong\ & [r+s]_*\big(\mathscr{L}_\psi(\beta(1/t')) \otimes (\mathrm{inv}^*\mathscr{K})|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big) \end{aligned}$$

This prove the general case of Theorem 0.2.

Frow now on, we assume $G(\alpha(t), r) = \{1\}$. By Lemma 2.10 (iii), we have $G(\beta(1/t'), r+s) = \{1\}$. If $\mathscr{K} = \mathscr{K}_\chi$ for some character

$$\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*,$$

then $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)$ is irreducible by Lemma 2.6, and hence $\mathscr{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big)$ is also irreducible by [18] 2.4.3 (i) c) and (ii) a). By Lemma 2.9, $\mathscr{L}_\psi(\beta(1/t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}$ is a direct factor of $[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big)$. For each $\mu \in \mu_{r+s}(k)$, $\beta(1/\mu t')$ is a solution of the system of equations in Theorem 0.2. So $\mathscr{L}_\psi(\beta(1/\mu t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}$ is also a direct factor of $[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big)$. As $G(\beta(1/t), r+s) = \{1\}$, the rank 1 representations of $\mathrm{Gal}(\bar{\eta}_{\infty'}/\eta_{\infty'})$ corresponding to $\mathscr{L}_\psi(\beta(1/\mu t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}$ are disjoint as μ go over $\mu_{r+s}(k)$. By Lemma 2.1 (ii), $[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big)$ is semisimple. Moreover, the rank of $[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big)$ is $r+s$ by [18] 2.4.3 (i) b). So we have

$$[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big) \cong \bigoplus_{\mu\in\mu_{r+s}(k)} \mathscr{L}_\psi(\beta(1/\mu t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}.$$

This is the isotypical decomposition of $[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big)$. The projection

$$[r+s]^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big) \to \mathscr{L}_\psi(\beta(1/t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}$$

defines a homomorphism

$$\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big) \to [r+s]_*\big(\mathscr{L}_\psi(\beta(1/t')) \otimes \mathscr{K}_{\chi^{-1}}|_{\eta_{\infty'}} \otimes ([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big).$$

If we apply $[r+s]^*$ to this homomorphism, we get an isomorphism. So this homomorphism itself is an isomorphism. This prove Theorem 0.2 in the case where $\mathscr{K} \cong \mathscr{K}_\chi$.

By Lemma 2.5, to prove Theorem 0.2 in general, we may assume

$$\mathscr{K} = \mathscr{K}_\chi \otimes U(n)$$

for some character $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$ and some integer n. Since $G(\alpha(t), r) = \{1\}$, the sheaf $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}|_{\eta_0}\big)$ is indecomposable by Lemma 2.6, and it contains $[r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)$ as an irreducible subsheaf. By [18] 2.4.3 (i) c) and (ii) a), $\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}|_{\eta_0}\big)\Big)$ is indecomposable, and it contains $\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K}_\chi|_{\eta_0}\big)\Big)$ as an irreducible subsheaf. So

$[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t'))\otimes(\mathrm{inv}^*\mathscr{K}_\chi)|_{\eta_{\infty'}}\otimes([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big)$ is an irreducible subsheaf of the indecomposable sheaf $\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}|_{\eta_0}\big)\Big)$. By Corollary 2.3 (i), we have

$$\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}|_{\eta_0}\big)\Big)\cong[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t'))\otimes(\mathrm{inv}^*\mathscr{K}_\chi)|_{\eta_{\infty'}}\otimes([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big)\otimes U(n')$$

for some integer n'. By [18] 2.4.3 (i) b), the rank of $\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}|_{\eta_0}\big)\Big)$ is

$$r\,\mathrm{rank}(\mathscr{K})+s\,\mathrm{rank}(\mathscr{K})=n(r+s),$$

whereas the rank of $[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t'))\otimes(\mathrm{inv}^*\mathscr{K}_\chi)|_{\eta_{\infty'}}\otimes([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big)\otimes U(n')$ is $n'(r+s)$. So we must have $n'=n$, and

$$\begin{aligned}
&\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}|_{\eta_0}\big)\Big)\\
\cong\ &[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t'))\otimes(\mathrm{inv}^*\mathscr{K}_\chi)|_{\eta_{\infty'}}\otimes([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big)\otimes U(n)\\
\cong\ &[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t'))\otimes((\mathrm{inv}^*\mathscr{K}_\chi)\otimes[r+s]^*U(n))|_{\eta_{\infty'}}\otimes([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big)\\
\cong\ &[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t'))\otimes(\mathrm{inv}^*(\mathscr{K}_\chi\otimes U(n)))|_{\eta_{\infty'}}\otimes([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big)\\
\cong\ &[r+s]_*\big(\mathscr{L}_\psi(\beta(1/t'))\otimes(\mathrm{inv}^*\mathscr{K})|_{\eta_{\infty'}}\otimes([s]^*\mathscr{K}_{\chi_2})|_{\eta_{\infty'}}\big).
\end{aligned}$$

This verifies the formula in Theorem 0.2. □

Proof of Theorem 0.1 (i). Here k is not necessarily algebraically closed. Notation as in Theorem 0.1 (i). By Lemma 1.10 (i), we have a projection

$$\delta^*\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)|_{\eta_0}\Big)\to\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}\Big(\frac{1}{2}s(r+s)a_{-s}t'^s\Big)\otimes G(\chi_2,\psi)\Big)|_{\eta_{0'}}.$$

It defines a homomorphism

$$\mathfrak{F}^{(0,\infty')}\Big([r]_*\big(\mathscr{L}_\psi(\alpha(t))\otimes\mathscr{K}\big)|_{\eta_0}\Big)\to\Big(\delta_*\Big(\mathscr{L}_\psi(\beta(t'))\otimes\mathscr{K}\otimes\mathscr{K}_{\chi_2}\Big(\frac{1}{2}s(r+s)a_{-s}t'^s\Big)\otimes G(\chi_2,\psi)\Big)\Big)|_{\eta_{\infty'}}.$$

To prove it is an isomorphism, we may make base extension from k to its algebraic closure $\bar{k}$. Over algebraically closed field, it is an isomorphism by the above proof of the $\mathscr{K}=\mathscr{K}_\chi$ case of Theorem 0.2. □

Proof of Theorems 0.1 (ii)-(iii), 0.3 and 0.4. The proof of Theorems 0.1 (ii) and 0.3 for the case $s>r$, and the proof of Theorems 0.1 (iii) and 0.4 for the case $s<r$ are similar to the above proof of Theorems 0.1 (i) and 0.2 by using the same group theoretical argument and Lemma 1.10 (ii)-(iii).

In the notation of Theorem 0.3, if $s < r$, then the slopes of $[r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}\big)|_{\eta_\infty}$ are $\frac{s}{r} \leq 1$. (Confer [17] 1.13.) By [18] 2.4.3 (iii) b), we then have

$$\mathfrak{F}^{(\infty,\infty')}\left([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}\big)|_{\eta_\infty}\right) = 0.$$

In the notation of Theorem 0.4, if $s > r$, then the slopes of $[r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}\big)|_{\eta_\infty}$ are $\frac{s}{r} \geq 1$. By [18] 2.4.3 (iii) b), we then have

$$\mathfrak{F}^{(\infty,0')}\left([r]_*\big(\mathscr{L}_\psi(\alpha(1/t)) \otimes \mathscr{K}\big)|_{\eta_\infty}\right) = 0.$$

Similarly we can prove Theorem 0.1 (ii) for the case $s < r$ and Theorem 0.1 (iii) for the case $s > r$. □

Proof of Proposition 0.5. By (b), there exists a finite Galois extension L of $k((t))$ such that ρ factors through $I_0 = \mathrm{Gal}(L/k((t)))$. The maximal tamely ramified extension of $k((t))$ contained in L is of the form $k((T))$, where $T = \sqrt[N]{t}$ is an N-th root of t for some positive integer N prime to p. Let $P_0 = \mathrm{Gal}(L/k((T)))$. $\mathrm{Res}^{I_0}_{P_0}(V)$ is semisimple. By (a), $\mathrm{Res}^{I_0}_{P_0}(V)$ factors through $P_0^{\mathrm{ab}} = P_0/[P_0, P_0]$. So $\mathrm{Res}^{I_0}_{P_0}(V)$ is a direct sum of one dimensional representations. Also by (a), the image of P_0 under each of these one dimensional representations is contained in

$$\mu_p(\overline{\mathbb{Q}}_\ell) = \{\mu \in \overline{\mathbb{Q}}_\ell^* | \mu^p = 1\}.$$

The additive character ψ is an isomorphism from $\mathbb{Z}/p$ to $\mu_p(\overline{\mathbb{Q}}_\ell)$. By the Artin-Schreier theory, each one dimensional direct summand of $\mathrm{Res}^{I_0}_{P_0}(V)$ is an Artin-Schreier representation, that is, a representation of P_0 corresponding to a sheaf $\mathscr{L}_\psi(\alpha(T))$ on $\mathrm{Spec}\, k((T))$ for some Laurent series $\alpha(T) \in k((T))$. Let $\{\alpha_1, \ldots, \alpha_m\}$ be a set of $\alpha_i(T) \in k((T))$ so that the representations of P_0 corresponding to the sheaves $\mathscr{L}_\psi(\alpha_i(T))$ on $\mathrm{Spec}\, k((T))$ give all one dimensional sub-representations of $\mathrm{Res}^{I_0}_{P_0}(V)$, and so that $\mathscr{L}_\psi(\alpha_i(T)) \ncong \mathscr{L}_\psi(\alpha_j(T))$ for $i \neq j$. Moreover, we can choose $\alpha_i(T)$ so that in their formal power series expansion, only non-positive powers of T are involved. Let

$$V = \bigoplus_{i=1}^{m} V_i$$

be the isotypical decomposition of $\mathrm{Res}^{I_0}_{P_0}(V)$ so that for each i, V_i is a direct sum of several copies of the representation corresponding to the sheaf $\mathscr{L}_\psi(\alpha_i(T))$. As in the beginning of the proof of Lemma 2.2 (i), I_0 acts on the set $\{V_1, \ldots, V_m\}$, and if I_1 is the stabilizer of V_1 under this action, then $P_0 \subset I_1$, V_1 is a representation of I_1, and $V = \mathrm{Ind}^{I_0}_{I_1}(V_1)$. The field L^{I_1} is a subfield of $L^{P_0} = k((T))$ and we have $L^{I_1} = k((S))$, where $S = T^{\frac{N}{r}}$ is an r-th root of t for some positive integer r dividing

N. Choose $g \in I_1$ so that $\frac{g(T)}{T} = \mu$ is a primitive $\frac{N}{r}$-th root of unity. Let $\rho_{V_1} : P_0 \to \mathrm{GL}(V_1)$ be the homomorphism defined by the representation V_1 and let $\rho^g_{V_1} : P_0 \to \mathrm{GL}(V_1)$ be defined by $\rho^g_{V_1}(\sigma) = \rho_{V_1}(g^{-1}\sigma g)$ for any $\sigma \in P_0$. Let $\rho_{\alpha_1} : P_0 \to \overline{\mathbb{Q}}^*_\ell$ be the character corresponding to the sheaf $\mathscr{L}_\psi(\alpha_1(T))$ on $\operatorname{Spec} k((T))$, and let $\rho^g_{\alpha_1} : P_0 \to \overline{\mathbb{Q}}^*_\ell$ be the character defined by $\rho^g_{\alpha_1}(\sigma) = \rho_{\alpha_1}(g^{-1}\sigma g)$ for any $\sigma \in P_0$. Then the representation defined by ρ_{V_1} (resp. $\rho^g_{V_1}$) is a direct sum of several copies of the irreducible representation defined by ρ_{α_1} (resp. $\rho^g_{\alpha_1}$). But $g : V_1 \to V_1$ is an isomorphism from the representation $\rho^g_{V_1}$ to the representation ρ_{V_1}. It follows that the representations defined by ρ_{α_1} and $\rho^g_{\alpha_1}$ are isomorphic. As ρ_{α_1} (resp. $\rho^g_{\alpha_1}$) corresponds to the sheaf $\mathscr{L}_\psi(\alpha_1(T))$ (resp. $\mathscr{L}_\psi(\alpha_1(\mu T))$) on $\operatorname{Spec} k((T))$, it follows that $\mathscr{L}_\psi(\alpha_1(T)) \cong \mathscr{L}_\psi(\alpha_1(\mu T))$. So there exists $\gamma(T) \in k((T))$ such that

$$\alpha_1(\mu T) - \alpha_1(T) = \gamma(T)^p - \gamma(T).$$

Write

$$\alpha_1(T) = \frac{a_{-s_1}}{T^{s_1}} + \frac{a_{-(s_1-1)}}{T^{s_1-1}} + \cdots + \frac{a_{-1}}{T}$$

with $a_{-s_1} \neq 0$. Denote by $s(-)$ the Swan conductor of a Galois representation. We have

$$s_1 = s(\mathscr{L}_\psi(\alpha_1(T))) \leq s(V_1) = s(\mathrm{Ind}_{I_1}^{I_0}(V_1)) = s(V) = s.$$

By (c), we have $s_1 < p$. The same argument as in the proof of Lemma 2.10 (i) shows that $\gamma(T)$ must be a formal power series, and $a_i = 0$ whenever $\frac{N}{r} \nmid i$. Let

$$\alpha(S) = \sum_{\frac{N}{r}|i} a_i S^{\frac{ir}{N}}.$$

Then we have $\alpha_1(T) = \alpha(T^{\frac{N}{r}})$. Denote the rank one representation of I_1 corresponding to the lisse $\overline{\mathbb{Q}}_\ell$-sheaf $\mathscr{L}_\psi(\alpha(S))$ on $\operatorname{Spec} k((S))$ also by $\mathscr{L}_\psi(\alpha(S))$. Then $\mathrm{Res}^{I_1}_{P_0}(\mathscr{L}_\psi(\alpha(S)))$ is the representation of P_0 corresponding to the sheaf $\mathscr{L}_\psi(\alpha_1(T))$ on $\operatorname{Spec} k((T))$. It follows that the representation $\mathrm{Res}^{I_1}_{P_0}(V_1 \otimes \mathscr{L}_\psi(\alpha(S))^{-1})$ is a direct sum of the trivial representation. Let $\mathscr{K} = V_1 \otimes \mathscr{L}_\psi(\alpha(S))^{-1}$. Then $\mathscr{K}$ is a tamely ramified representation of I_1. We have

$$V = \mathrm{Ind}_{I_1}^{I_0}(V_1) = \mathrm{Ind}_{I_1}^{I_0}(\mathscr{L}_\psi(\alpha(S)) \otimes \mathscr{K}).$$

As V is irreducible, $\mathscr{L}_\psi(\alpha(S)) \otimes \mathscr{K}$ is irreducible, and hence $\mathscr{K}$ is irreducible. So $\mathscr{K} = \mathscr{K}_\chi$ for some character $\chi : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}^*_\ell$ of finite order. □

Remark 2.11. If the statement in Remark 2.4 holds, then by Corollary 2.3 (i) and Proposition 0.5, the following generalization of Proposition 0.5 holds: Let $\rho : I \to \mathrm{GL}(V)$ be an indecomposable $\overline{\mathbb{Q}}_\ell$-representation. Suppose $\rho(P^p[P,P]) = 1$ and suppose the Swan conductor of ρ is less than p. Then V is isomorphic to the representation corresponding to the lisse $\overline{\mathbb{Q}}_\ell$-sheaf $[r]_*(\mathscr{L}_\psi(\alpha(t)) \otimes \mathscr{K})$ on $\eta_0 = \mathrm{Spec}\, k((t))$ for some positive integer r prime to p, some Laurent series $\alpha(t)$ and some tamely ramified sheaf $\mathscr{K}$ on η_0.

3 Local monodromy of hypergeometric sheaves

Let k be an algebraically closed field of characteristic p, and let

$$\lambda_1, \ldots, \lambda_n, \rho_1, \ldots, \rho_m : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$$

be characters. In [15] 8.2, Katz defines the hypergeometric complex $\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)$ in $D_c^b(\mathbb{G}_{m,k}, \overline{\mathbb{Q}}_\ell)$ using the multiplicative convolution. Let $j : \mathbb{G}_{m,k} \to \mathbb{A}^1_k$ be the canonical open immersion, and let $\mathrm{inv} : \mathbb{G}_{m,k} \to \mathbb{G}_{m,k}$ be the morphism defined by $t \mapsto \frac{1}{t}$. Recall the following properties of the hypergeometric complex ([15] 8.2.2 (2), 8.2.4, 8.2.5, 8.1.12).

(a) If $n = 1$ and $m = 0$, we have

$$\mathrm{Hyp}(!, \psi; \lambda_1; \emptyset) \cong j^*\mathscr{L}_\psi \otimes \mathscr{K}_{\lambda_1}[1].$$

(b) For any character $\lambda : \varprojlim_{(N,p)=1} \mu_N(k) \to \overline{\mathbb{Q}}_\ell^*$, we have

$$\mathrm{Hyp}(!, \psi; \lambda\lambda_1, \ldots, \lambda\lambda_n; \lambda\rho_1, \ldots, \lambda\rho_m) \cong \mathscr{K}_\lambda \otimes \mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m).$$

(c) In the case where $\lambda_n = 1$, we have

$$\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_{n-1}, 1; \rho_1, \ldots, \rho_m) \cong j^*\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_{n-1}; \rho_1, \ldots, \rho_m)).$$

(d) We have

$$\mathrm{inv}^*\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m) \cong \mathrm{Hyp}(!, \psi^{-1}; \rho_1^{-1}, \ldots, \rho_m^{-1}; \lambda_1^{-1}, \ldots, \lambda_n^{-1}).$$

The above properties can also be used to define $\mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m)$. Indeed, by properties (b) and (c), we have

$$\begin{aligned} & \mathrm{Hyp}(!, \psi; \lambda_1, \ldots, \lambda_n; \rho_1, \ldots, \rho_m) \\ \cong\ & \mathscr{K}_{\lambda_n} \otimes \mathrm{Hyp}(!, \psi; \lambda_1\lambda_n^{-1}, \ldots, \lambda_{n-1}\lambda_n^{-1}, 1; \rho_1\lambda_n^{-1}, \ldots, \rho_m\lambda_n^{-1}) \\ \cong\ & \mathscr{K}_{\lambda_n} \otimes j^*\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!, \psi; \lambda_1\lambda_n^{-1}, \ldots, \lambda_{n-1}\lambda_n^{-1}; \rho_1\lambda_n^{-1}, \ldots, \rho_m\lambda_n^{-1})). \end{aligned}$$

This reduces the definition of $\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_n;\rho_1,\ldots,\rho_m)$ to the case where $n=0$. By properties (d), (b) and (c), for the case where $n=0$, we have

$$\begin{aligned}
& \mathrm{Hyp}(!,\psi;\emptyset;\rho_1,\ldots,\rho_m)\\
\cong\ & \mathrm{inv}^*\mathrm{Hyp}(!,\psi^{-1};\rho_1^{-1},\ldots,\rho_m^{-1};\emptyset)\\
\cong\ & \mathrm{inv}^*(\mathscr{K}_{\rho_m^{-1}}\otimes\mathrm{Hyp}(!,\psi^{-1};\rho_1^{-1}\rho_m,\ldots,\rho_{m-1}^{-1}\rho_m,1;\emptyset))\\
\cong\ & \mathrm{inv}^*(\mathscr{K}_{\rho_m^{-1}}\otimes j^*\mathfrak{F}_{\psi^{-1}}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi^{-1};\rho_1^{-1}\rho_m,\ldots,\rho_{m-1}^{-1}\rho_m;\emptyset))),
\end{aligned}$$

where $\mathfrak{F}_{\psi^{-1}}$ denotes the Fourier transformation defined by the additive character ψ^{-1}. This reduces the definition of $\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_n;\rho_1,\ldots,\rho_m)$ to that in the case where $n=1$ and $m=0$, for which we can use the property (a).

To prove Proposition 0.7 in the Introduction, we first prove Proposition 0.8.

Proof of Proposition 0.8. When $n=1$, our assertion follows from the fact that

$$\mathrm{Hyp}(!,\psi;\lambda_1;\emptyset)[-1]\cong j^*\mathscr{L}_\psi\otimes\mathscr{K}_{\lambda_1}.$$

Suppose the lemma holds if there are $n-1$ characters λ's. To prove the lemma for n characters λ's, we use the fact that

$$\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_n;\emptyset)\cong\mathscr{K}_{\lambda_n}\otimes\mathrm{Hyp}(!,\psi;\lambda_1\lambda_n^{-1},\ldots,\lambda_{n-1}\lambda_n^{-1},1;\emptyset)$$

to reduce to the case where $\lambda_n=1$. We have

$$\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)=j^*\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)).$$

By the induction hypothesis, we have

$$\big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_\infty}\cong[n-1]_*\big(\mathscr{L}_\psi((n-1)t)\otimes\mathscr{K}_{\lambda_1\cdots\lambda_{n-1}}\otimes\mathscr{K}_{\chi_2{}^{n-2}}\big)|_{\eta_\infty}.$$

So

$$\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_n;\emptyset)[-1]\big)|_{\eta_0}\cong[n-1]_*\left(\mathscr{L}_\psi\left(\frac{n-1}{t}\right)\otimes\mathrm{inv}^*(\mathscr{K}_{\lambda_1\cdots\lambda_{n-1}}\otimes\mathscr{K}_{\chi_2{}^{n-2}})\right)|_{\eta_0}.\quad(1)$$

By the induction hypothesis, $\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]$ is an irreducible lisse sheaf of rank $n-1$ on $\mathbb{G}_{m,k}$. Combined with (1), we get

$$j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)\cong j_{!*}\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset),$$

and $j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)$ is an irreducible perverse sheaf of type (T_3) in [18] 1.4.2. So $\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset))$ is an irreducible perverse sheaf on $\mathbb{A}^1_k$ of type (T_3) by [18] 1.4.2.1 (ii). By the induction hypothesis, we have

$$\begin{array}{rcl}\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_\infty} & \cong & \mathrm{inv}^*\big(\big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_0}\big)\\ & \cong & \mathrm{inv}^*\big(\bigoplus_\lambda\big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0}\big)\\ & \cong & \bigoplus_\lambda\big(\mathrm{inv}^*(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda)))\big)|_{\eta_\infty},\end{array}\tag{2}$$

where $\mathrm{mult}_0(\lambda)$ denotes the number of times that λ appears in $\lambda_1,\ldots,\lambda_{n-1}$. By (2) and [18] 2.3.1.3 (i), $\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset))$ is lisse on $\mathbb{G}_{m,k}$. It follows that $\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)[-1]$ is an irreducible lisse sheaf on $\mathbb{G}_{m,k}$, and we have

$$\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset))=j_{!*}\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset),$$

and the long exact sequence in [18] 2.3.2 gives rise to a short exact sequence

$$\begin{array}{rcl}0 & \to & \big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)[-1]\big)_{\bar\eta_0}^{\mathrm{Gal}(\bar\eta_0/\eta_0)}\\ & \to & \big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)[-1]\big)_{\bar\eta_0}\\ & \to & \mathfrak{F}^{(\infty,0)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_\infty}\big)\ \to\ 0.\end{array}\tag{3}$$

We have

$$\begin{array}{rcl} & & \mathfrak{F}^{(\infty,0)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_\infty}\big)\\ & \cong & \mathfrak{F}^{(\infty,0)}\big(\bigoplus_\lambda\big(\mathrm{inv}^*(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda)))\big)|_{\eta_\infty}\big)\\ & \cong & \bigoplus_\lambda\big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0}.\end{array}\tag{4}$$

Using the formula in [18] 2.3.1.1 (i) and (iii), one can check

$$\mathrm{rank}\big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)[-1]\big)_{\bar\eta_0}^{\mathrm{Gal}(\bar\eta_0/\eta_0)}=1,\tag{5}$$

$$\mathrm{rank}\big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)[-1]\big)=n.\tag{6}$$

It follows from (3)-(6) that

$$\big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)[-1]\big)|_{\eta_0}\cong U(\mathrm{mult}_0(1))|_{\eta_0}\bigoplus\bigoplus_{\lambda\neq1}\big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0},$$

where $\mathrm{mult}_0(1)$ is the number of trivial characters in $\lambda_1,\ldots,\lambda_{n-1},1$.

By [18] 2.3.3.1 (iii), we have

$$\begin{array}{rcl}\big(\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1},1;\emptyset)[-1]\big)|_{\eta_\infty} & \cong & \mathfrak{F}^{(0,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_0}\big)\\ & & \bigoplus\mathscr{F}^{(\infty,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_\infty}\big).\end{array}$$

By (2) and [18] 2.4.3 (iii) b), we have

$$\mathfrak{F}^{(\infty,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\ldots,\lambda_{n-1};\emptyset)[-1]\big)|_{\eta_\infty}\big)=0.$$

Using the formula in Theorem 0.2 and (1), we find

$$\mathfrak{F}^{(0,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\emptyset)[-1]\big)|_{\eta_0}\big) \cong [n]_*\big(\mathscr{L}_\psi(nt)\otimes\mathscr{K}_{\lambda_1\cdots\lambda_{n-1}}\otimes\mathscr{K}_{\chi_2{}^{n-1}}\big)|_{\eta_\infty}.$$

It follows that

$$\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\emptyset)[-1]\big)|_{\eta_\infty} \cong [n]_*\big(\mathscr{L}_\psi(nt)\otimes\mathscr{K}_{\lambda_1\cdots\lambda_{n-1}}\otimes\mathscr{K}_{\chi_2{}^{n-1}}\big)|_{\eta_\infty}.$$

□

Using Proposition 0.8 and the fact that

$$\mathrm{Hyp}(!,\psi;\emptyset;\rho_1,\dots,\rho_m) \cong \mathrm{inv}^*\mathrm{Hyp}(!,\psi^{-1};\rho_1^{-1},\dots,\rho_m^{-1};\emptyset),$$

we get the following.

Corollary 3.1. *Suppose p is relatively prime to $2,3,\dots,m$. Then* $\mathrm{Hyp}(!,\psi;\emptyset;\rho_1,\dots,\rho_m)[-1]$ *is an irreducible lisse sheaf of rank m on $\mathbb{G}_{m,k}$, and we have*

$$\begin{aligned}
\big(\mathrm{Hyp}(!,\psi;\emptyset;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty} &\cong \bigoplus_\rho\big(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty},\\
\big(\mathrm{Hyp}(!,\psi;\emptyset;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_0} &\cong [m]_*\left(\mathscr{L}_\psi\left(-\frac{m}{t}\right)\otimes\mathscr{K}_{\rho_1\cdots\rho_m}\otimes\mathscr{K}_{\chi_2{}^{m-1}}\right)|_{\eta_0}.
\end{aligned}$$

We now prove Proposition 0.7.

Proof of Proposition 0.7. Using the fact that

$$\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m) \cong \mathrm{inv}^*\mathrm{Hyp}(!,\psi^{-1};\rho_1^{-1},\dots,\rho_m^{-1};\lambda_1^{-1},\dots,\lambda_n^{-1}),$$

we can deduce (i) from (ii). Let's prove (ii). When $n=0$, this follows from Corollary 3.1. Suppose $1\le n<m$ and suppose the lemma holds if there are $n-1$ characters λ's. To prove the lemma for n characters λ's, we use the fact that

$$\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m) \cong \mathscr{K}_{\lambda_n}\otimes\mathrm{Hyp}(!,\psi;\lambda_1\lambda_n^{-1},\dots,\lambda_{n-1}\lambda_n^{-1},1;\rho_1\lambda_n^{-1},\dots,\rho_m\lambda_n^{-1})$$

to reduce to the case where $\lambda_n=1$. Then by the disjointness of λ's and ρ's, we have $\rho_i\neq 1$ for all $1\le i\le m$. We have

$$\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m) = j^*\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)).$$

By the induction hypothesis, $\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]$ is an irreducible lisse sheaf of rank m on $\mathbb{G}_{m,k}$, and

$$\big(\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_0}\cong\bigoplus_{\rho}\big(\mathrm{inv}^*(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho)))\big)|_{\eta_0}.$$

Since $\rho_i\neq 1$ for all $1\le i\le m$, this implies that

$$j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)\cong j_{!*}\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m),$$

and $j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)$ is an irreducible perverse sheaf of type (T_3) in [18] 1.4.2. So $\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m))$ is an irreducible perverse sheaf on $\mathbb{A}^1_k$ of type (T_3) by [18] 1.4.2.1 (ii). By the induction hypothesis, we have

$$\begin{aligned}
&\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}\\
\cong\ &\mathrm{inv}^*\Big([m-(n-1)]_*\Big(\mathscr{L}_\psi\Big(-\tfrac{m-(n-1)}{t}\Big)\otimes\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_{n-1}^{-1}\rho_1\cdots\rho_m}\otimes\mathscr{K}_{{\chi_2}^{n+m-2}}\Big)\\
&\qquad\bigoplus\bigoplus\nolimits_\lambda\big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)\Big)|_{\eta_\infty}\qquad(7)\\
\cong\ &[m-(n-1)]_*\Big(\mathscr{L}_\psi(-(m-(n-1))t)\otimes\mathrm{inv}^*\big(\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_{n-1}^{-1}\rho_1\cdots\rho_m}\otimes\mathscr{K}_{{\chi_2}^{n+m-2}}\big)\Big)|_{\eta_\infty}\\
&\quad\bigoplus\bigoplus\nolimits_\lambda\big(\mathrm{inv}^*(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda)))\big)|_{\eta_\infty}.
\end{aligned}$$

So all the breaks of $\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}$ are <1. By [18] 2.3.1.3 (i), $\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m))$ is lisse on $\mathbb{G}_{m,k}$. It follows that $\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m)[-1]$ is an irreducible lisse sheaf on $\mathbb{G}_{m,k}$, we have

$$\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m))\cong j_{!*}\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m),$$

and the long exact sequence in [18] 2.3.2 gives rise to the short exact sequence

$$\begin{aligned}
0\ &\to\ \big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m)[-1]\big)_{\bar\eta_0}^{\mathrm{Gal}(\bar\eta_0/\eta_0)}\\
&\to\ \big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m)[-1]\big)_{\bar\eta_0}\qquad(8)\\
&\to\ \mathfrak{F}^{(\infty,0)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}\big)\ \to\ 0.
\end{aligned}$$

Using the formulas in [18] 2.3.1.1 (i) and (iii), one can check

$$\mathrm{rank}\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m)[-1]\big)_{\bar\eta_0}^{\mathrm{Gal}(\bar\eta_0/\eta_0)}=1,\qquad(9)$$

$$\mathrm{rank}\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m)[-1]\big)=m.\qquad(10)$$

Using the formula in Theorem 0.4 and (7), we find that

$$\begin{aligned}
&\mathfrak{F}^{(\infty,0)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}\big)\\
\cong\ &[m-n]_*\Big(\mathscr{L}_\psi\big(-\tfrac{m-n}{t}\big)\otimes\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_n^{-1}\rho_1\cdots\rho_m}\otimes\mathscr{K}_{{\chi_2}^{n+m-1}}\Big)|_{\eta_0}\bigoplus\bigoplus\nolimits_\lambda\big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0}.\qquad(11)
\end{aligned}$$

It follows from (8)-(11) that

$$\begin{aligned}
&\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_0}\\
\cong\ & [m-n]_*\left(\mathscr{L}_\psi\left(-\frac{m-n}{t}\right)\otimes\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_n^{-1}\rho_1\cdots\rho_m}\otimes\mathscr{K}_{\chi_2{}^{n+m-1}}\right)|_{\eta_0}\\
&\bigoplus U(\mathrm{mult}_0(1))|_{\eta_0}\bigoplus\bigoplus_{\lambda\neq 1}\big(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda))\big)|_{\eta_0}.
\end{aligned}$$

By [18] 2.3.3.1 (iii), we have

$$\begin{aligned}
&\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}\\
\cong\ & \mathfrak{F}^{(0,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_0}\big)\\
&\bigoplus\mathscr{F}^{(\infty,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_n,\dots,\rho_m)[-1]\big)|_{\eta_\infty}\big).
\end{aligned}$$

Since $\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}$ has breaks <1, we have

$$\mathfrak{F}^{(\infty,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}\big)=0$$

by [18] 2.4.3 (iii) b). We have

$$\begin{aligned}
&\mathfrak{F}^{(0,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_0}\big)\\
\cong\ & \mathfrak{F}^{(0,\infty)}\big(\bigoplus_\rho\big(\mathrm{inv}^*(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho)))\big)|_{\eta_0}\big)\\
\cong\ & \bigoplus_\rho\big(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty}.
\end{aligned}$$

It follows that

$$\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_m)[-1]\big)|_{\eta_\infty}\cong\bigoplus_\rho\big(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty}.$$

Finally, we prove (iii). First consider the case $\lambda_n=1$. By the disjointness of λ's and ρ's, we have $\rho_i\neq 1$ for all i. We have

$$\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_n)\cong j^*\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)).$$

By (ii), $\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]$ is an irreducible lisse sheaf on $\mathbb{G}_{m,k}$ and

$$\big(\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_0}\cong\bigoplus_\rho\big(\mathrm{inv}^*(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho)))\big)|_{\eta_0}.\tag{12}$$

Since $\rho_i\neq 1$ for all i, this implies that

$$j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)\cong j_{!*}\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n),$$

and $j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)$ is an irreducible perverse sheaf of type (T_3) in [18] 1.4.2. So $\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n))$ is an irreducible perverse sheaf on $\mathbb{A}^1_k$ of type (T_3) by [18] 1.4.2.1 (ii). By (ii), we have

$$\begin{array}{rl} & \big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_\infty} \\ \cong & \big(\mathscr{L}_\psi(-t)\otimes \mathrm{inv}^*\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_{n-1}^{-1}\rho_1\cdots\rho_n}\big)|_{\eta_\infty}\bigoplus\bigoplus_\lambda\big(\mathrm{inv}^*(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda)))\big)|_{\eta_\infty}. \end{array} \tag{13}$$

By (13) and [18] 2.3.1.2, $\mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n))$ is lisse on $\mathbb{G}_{m,k}-\{1\}$. Therefore $\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_n)[-1]$ is an irreducible lisse sheaf when restricted to $\mathbb{G}_{m,k}-\{1\}$. Using [18] 2.3.1.1 (i), one can show its rank is n.

Let $\mathrm{tran}:\mathbb{A}^1_k\to\mathbb{A}^1_k$ be the translation by 1. By the definition of the Fourier transformation, we have

$$\begin{array}{rl} & \mathfrak{F}(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)) \\ \cong & \mathrm{tran}^*\mathfrak{F}(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)). \end{array} \tag{14}$$

Note that $\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)$ is also an irreducible perverse sheaf on $\mathbb{A}^1_k$ of type (T_3). Hence $\mathfrak{F}(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n))$ is an irreducible perverse sheaf on $\mathbb{A}^1_k$ of type (T_3). So we have

$$\begin{array}{rl} & \mathscr{H}^{-1}\big(\mathfrak{F}(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n))\big)_{\bar 0} \\ \cong & \mathscr{H}^{-1}\big(\mathfrak{F}(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n))\big)_{\bar\eta_0}^{\mathrm{Gal}(\bar\eta_0/\eta_0)}, \end{array} \tag{15}$$

and the long exact sequence in [18] 2.3.2 gives rise to the short exact sequence

$$\begin{array}{rcl} 0 & \to & \mathscr{H}^{-1}\big(\mathfrak{F}(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n))\big)_{\bar 0} \\ & \to & \mathscr{H}^{-1}\big(\mathscr{F}(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n))\big)_{\bar\eta_0} \\ & \to & \mathfrak{F}^{(\infty,0)}\big(\big(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_\infty}\big) \quad\to\quad 0. \end{array} \tag{16}$$

We have

$$\begin{array}{rl} & \mathfrak{F}^{(\infty,0)}\big(\big(\mathscr{L}_\psi\otimes j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_\infty}\big) \\ \cong & \mathfrak{F}^{(\infty,0)}\big(\big(\mathrm{inv}^*\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_{n-1}^{-1}\rho_1\cdots\rho_n}\big)|_{\eta_\infty}\bigoplus\bigoplus_\lambda\big(\mathscr{L}_\psi\otimes\mathrm{inv}^*(\mathscr{K}_\lambda\otimes U(\mathrm{mult}_0(\lambda)))|_{\eta_\infty}\big) \\ \cong & \mathscr{K}_{\lambda_1^{-1}\cdots\lambda_{n-1}^{-1}\rho_1\cdots\rho_n}|_{\eta_0} \end{array} \tag{17}$$

by [18] 2.4.3 (ii) b) and 2.5.3.1. Applying tran^* to (15)-(17) and using (14), we see that

$$\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_n)[-1]\big)_{\bar 1}\cong\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_n)[-1]\big)_{\bar\eta_1}^{\mathrm{Gal}(\bar\eta_1/\eta_1)},$$

and we have a short exact sequence

$$\begin{array}{rcl} 0 & \to & \big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_n)[-1]\big)_{\bar\eta_1}^{\mathrm{Gal}(\bar\eta_1/\eta_1)} \\ & \to & \big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_n)[-1]\big)_{\bar\eta_1} \\ & \to & (\mathrm{tran}^*\mathscr{K}_{\lambda_1^{-1}\cdots\lambda_{n-1}^{-1}\rho_1\cdots\rho_n})|_{\bar\eta_1}\to 0. \end{array}$$

By (13), all the breaks of $(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1])|_{\eta_\infty}$ are ≤ 1. It follows that

$$\mathfrak{F}^{(\infty,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_\infty}\big)=0.$$

By (12) and [18] 2.5.3.1, we have

$$\mathfrak{F}^{(0,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_0}\big)\cong\bigoplus_\rho\big(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty}.$$

By [18] 2.3.3.1 (iii), we have

$$\begin{aligned}
&\big(\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1},1;\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_\infty}\\
\cong\ &\mathfrak{F}^{(0,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_1,\dots,\rho_n)[-1]\big)|_{\eta_0}\big)\\
&\qquad\bigoplus\mathscr{F}^{(\infty,\infty)}\big(\big(j_!\mathrm{inv}^*\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_{n-1};\rho_n,\dots,\rho_n)[-1]\big)|_{\eta_\infty}\big)\\
\cong\ &\bigoplus_\rho\big(\mathscr{K}_\rho\otimes U(\mathrm{mult}_\infty(\rho))\big)|_{\eta_\infty}.
\end{aligned}$$

This proves those statements in Proposition 0.7 (iii) about the behavior at 1 and at ∞ for $\lambda_n=1$. For general λ_n, we use the fact that

$$\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_n)\cong\mathscr{K}_{\lambda_n}\otimes\mathrm{Hyp}(!,\psi;\lambda_1\lambda_n^{-1},\dots,\lambda_{n-1}\lambda_n^{-1},1;\rho_1\lambda_n^{-1},\dots,\rho_n\lambda_n^{-1}).$$

For the statement in Proposition 0.7 (iii) about the behavior at 0, we use the fact that

$$\mathrm{Hyp}(!,\psi;\lambda_1,\dots,\lambda_n;\rho_1,\dots,\rho_n)\cong\mathrm{inv}^*\mathrm{Hyp}(!,\psi^{-1};\rho_1^{-1},\dots,\rho_n^{-1};\lambda_1^{-1},\dots,\lambda_n^{-1}).$$

□

References

[1] A. Abbes and T. Saito, Local Fourier Transform and Epsilon Factors, arXiv: math/0809.0180 (2008).

[2] V. I. Arnold, Mathematical methods of classical mechanics, Springer-Verlag (1989).

[3] M. Artin, A. Grothendieck, and J.-L. Verdier, Théorie des topos et cohomologie étale des schémas (SGA 4), *Lecture Notes in Math.* 269, 270, 305, Springer-Verlag (1972-1973).

[4] E. T. Copson, Asymptotic expansions, Cambridge University Press (1965).

[5] P. Deligne et al, Cohomologie étale (SGA $4\frac{1}{2}$), *Lecture Notes in Math.* 569, Springer-Verlag (1977).

[6] P. Deligne, Letters to the author, April 29 and May 6, 2008.

[7] P. Deligne, La conjecture de Weil II, *Publ. Math. IHES* 52 (1980), 137-252.

[8] J. Fang, Calculation of local Fourier transforms for formal connections, arXiv: math/0707.0090 (2007).

[9] L. Fu and D. Wan, L-functions for symmetric products of Kloosterman sums, *J. Reine Angew. Math.* 589 (2005), 79-103.

[10] A. Grothendieck, Revêtements Étales et Groupe Fondamental (SGA 1), *Lecture Notes in Math.* 224, Springer-Verlag (1971).

[11] A. Grothendieck, P. Deligne, and N. Katz, Groupes de monodromie en géométrie algébrique (SGA 7), *Lecture Notes in Math.* 228, 340, Springer-Verlag (1972-1973).

[12] A. Grothendieck and D. Dieudonné, Étude globale élémentaire de quelques classes de morphismes (EGA II), *Publ. Math. IHES* 8 (1961).

[13] A. Grothendieck and D. Dieudonné, Étude cohomologique des faisceaux cohérents (EGA III), *Publ. Math. IHES* 11 (1961), 17 (1963).

[14] A. Grothendieck and D. Dieudonné, Étude locale des schémas et des morphismes de schémas (EGA IV), *Publ. Math. IHES* 20 (1964), 24 (1965), 28 (1966), 32 (1967).

[15] N. Katz, Exponential sums and differential equations, *Annals of Mathematics Studies* 124, Princeton University Press (1990).

[16] N. Katz, Exponential sums over finite fields and differential equations over the complex numbers: some interactions, *Bull. Amer. Math. Soc.* 23 (1990), 269-309.

[17] N. Katz, Gauss sums, Kloosterman sums, and monodromy groups, *Annals of Mathematics Studies* 116, Princeton University Press (1988).

[18] G. Laumon, Transformation de Fourier, constantes d'équations fontionnelles, et conjecture de Weil, *Publ. Math. IHES* 65 (1987), 131-210.

[19] L. Ramero, On a class of étale analytic sheaves, *J. Alg. Geo.* 7 (1998), 405-504.

[20] C. Sabbah, An explicit stationary phase formula for the local formal Fourier-Laplace transform, Singularities I, 309-330, *Contemp. Math.*, 474, Amer. Math. Soc., Providence, RI, 2008.

[21] J.-P. Serre, Linear representations of finite groups, Springer-Verlag (1977).

[22] J.-P. Serre and J. Tate, Good reduction of abelian varieties, *Ann. of Math.* 88 (1968), 492-517.